

# INDIANA

## NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
Form originally created for use in New York. Indiana Form version date: November 1, 2010

Scientific name: Microstegium vimineum USDA Plants Code: MIVI  
 Common names: Japanese stiltgrass, Eulalia, Nepalese browntop  
 Native distribution: India and Asia, including Bhutan, China, Japan, Korea, Malaysia, Myanmar, Nepal, and Vietnam  
 Date assessed: July 16, 2012  
 Assessors: Pia Marie Paulone and Ellen Jacquart  
 Reviewers: Jason Larson and Roger Hedge  
 Date Approved: September 21, 2012

**Indiana Invasiveness Rank:** Very High (Relative Maximum Score >80.00)

Invasiveness Ranking Summary (see details under appropriate sub-section)		Total (Total Answered*) Possible	Total
1	Ecological impact	40 ( <u>40</u> )	33
2	Biological characteristic and dispersal ability	25 ( <u>25</u> )	20
3	Ecological amplitude and distribution	25 ( <u>25</u> )	24
4	Difficulty of control	10 ( <u>10</u> )	7
	Outcome score	100 ( <u>100</u> ) <sup>b</sup>	84 <sup>a</sup>
	Relative maximum score <sup>†</sup>		84.00
	Indiana Invasiveness Rank <sup>§</sup>	Very High	

\* For questions answered "unknown" do not include point value in "Total Answered Points Possible." If "Total Answered Points Possible" is less than 70.00 points, then the overall invasive rank should be listed as "Unknown."

<sup>†</sup>Calculated as 100(a/b) to two decimal places.

<sup>§</sup>Very High >80.00; High 70.00–80.00; Moderate 50.00–69.99; Low 40.00–49.99; Insignificant <40.00

### A. DISTRIBUTION (KNOWN/POTENTIAL):

<p>A1. Has this species been documented to persist without cultivation in IN? (reliable source; voucher not required)</p> <p><input checked="" type="checkbox"/> Yes – continue to A2.2</p> <p><input type="checkbox"/> No – continue to A2.1</p> <hr/> <p>A2. What is the likelihood that this species will occur and persist outside of cultivation given the climate in Indiana? (obtain from occurrence data in other states with similar climates)</p> <p><input type="checkbox"/> Likely – continue to A3</p> <p><input type="checkbox"/> Not likely</p>	<div style="border: 1px solid black; padding: 5px;"> <p><b>Legend</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> IPSAWG</li> <li><input checked="" type="checkbox"/> PLANTS</li> <li><input type="checkbox"/> CAPS</li> <li><input checked="" type="checkbox"/> EDDMaps</li> <li><input type="checkbox"/> NO RECORD</li> </ul> <p>Date: 9/18/2012</p> </div>
--	---

**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
Form originally created for use in New York. Indiana Form version date: November 1, 2010

**Documentation:**

Sources of information: Range maps compiled from PLANTS database, <http://plants.usda.gov/java/>; Indiana CAPS database, <http://extension.entm.purdue.edu/CAPS/index.html>; Indiana IPSAWG reports (unpublished); and EDDMapS reports, <http://eddmaps.org/>

A3 Describe the potential or known suitable habitats within Indiana (underlined). Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

<p><b>Aquatic Habitats</b></p> <p>Rivers/streams Natural lakes and ponds Reservoirs/impoundments*</p>	<p><b>Wetland Habitats</b></p> <p>Marshes Fens Bogs <u>Shrub swamps</u> <u>Forested wetlands/riparian</u> Beaches/dunes <u>Ditches*</u></p>	<p><b>Upland Habitats</b></p> <p><u>Forest</u> <u>Savannas</u> <u>Barrens</u> Prairies Cultivated* <u>Old Fields*</u> <u>Roadsides*</u></p>
---	---	---

Other potential or known suitable habitats within Indiana:

No additional habitats.

**Documentation:**

Sources of information:

Gleason & Cronquist, 1991; Swearingen & Adams, 2007; Brooklyn Botanic Garden, 2008; U.S.D.A., 2008.

**B. INVASIVENESS RANKING**

**Questions apply to areas similar in climate and habitats to Indiana unless specified otherwise.**

*1. ECOLOGICAL IMPACT*

1.1. Impact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire regime, geomorphological changes (erosion, sedimentation rates), hydrologic regime, nutrient and mineral dynamics, light availability, salinity, pH)

- |  |                                      |
|--|--------------------------------------|
| <p>A. No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (&gt;10 occurrences in minimally managed areas), has been well-studied (&gt;10 reports/publications), and has been present in the northeast for &gt;100 years.</p> <p>B. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability)</p> <p>C. Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl)</p> <p>D. Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species)</p> <p>U. Unknown</p> | <p>0</p> <p>3</p> <p>7</p> <p>10</p> |
|--|--------------------------------------|

Score 

10
----

**Documentation:**

Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)

Increases soil pH, increases fire intensity in forests.

Sources of information:

Ehrenfeld et al., 2001; Kourtev et al., 2002; Swearingen & Adams, 2007; Flory, 2011

1.2. Impact on Natural Community Structure

**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
Form originally created for use in New York. Indiana Form version date: November 1, 2010

---

- A. No perceived impact; establishes in an existing layer without influencing its structure 0
- B. Influences structure in one layer (e.g., changes the density of one layer) 3
- C. Significant impact in at least one layer (e.g., creation of a new layer or elimination of an existing layer) 7
- D. Major alteration of structure (e.g., covers canopy, eradicating most or all layers below) 10
- U. Unknown

Score

**Documentation:**

Identify type of impact or alteration:

Oftentimes will substantially increase herb layer. Since *M. vimineum* is less palatable to white-tailed deer than are native plants, deer browse pressure may be increased on native plants which may reduce their recruitment (Eschtruth & Battles, 2008). It increases fire intensity in forests, causing mortality in tree seedlings.

Sources of information:

Winter et al., 1982; Redman, 1995; La Fleur, 1996; Horton & Neufeld, 1998; Claridge & Franklin, 2003; Cole & Weltzin, 2004, 2005; Flory, 2011.

**1.3. Impact on Natural Community Composition**

- A. No perceived impact; causes no apparent change in native populations 0
- B. Influences community composition (e.g., reduces the number of individuals in one or more native species in the community) 3
- C. Significantly alters community composition (e.g., produces a significant reduction in the population size of one or more native species in the community) 7
- D. Causes major alteration in community composition (e.g., results in the extirpation of one or several native species, reducing biodiversity or change the community composition towards species exotic to the natural community) 10
- U. Unknown

Score

**Documentation:**

Identify type of impact or alteration:

Oftentimes will form a monoculture in the herb layer. Low palatability to white-tailed deer may result in increased deer browse on native plant species, which may reduce their recruitment (Eschtruth & Battles, 2008). Increases fire intensity, causing mortality in tree seedling layer.

Sources of information:

Winter et al., 1982; Redman, 1995; La Fleur, 1996; Horton & Neufeld, 1998; Claridge & Franklin, 2003; Cole & Weltzin 2004, 2005; Flory; 2011.

**1.4. Impact on other species or species groups (cumulative impact of this species on the animals, fungi, microbes, and other organisms in the community it invades.**

Examples include reduction in nesting/foraging sites; reduction in habitat connectivity; injurious components such as spines, thorns, burrs, toxins; suppresses soil/sediment microflora; interferes with native pollinators and/or pollination of a native species; hybridizes with a native species; hosts a non-native disease which impacts a native species)

- A. Negligible perceived impact 0
- B. Minor impact 3
- C. Moderate impact 7
- D. Severe impact on other species or species groups 10
- U. Unknown

Score

**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
Form originally created for use in New York. Indiana Form version date: November 1, 2010

**Documentation:**

Identify type of impact or alteration:  
Dominance of herb layer and alteration of soil chemistry.

Sources of information:

Winter et al., 1982; Redman, 1995; La Fleur, 1996; Horton & Neufeld, 1998; Ehrenfeld et al., 2001; Claridge & Franklin, 2003; Cole & Weltzin 2004, 2005; Swearingen & Adams, 2007.

Total Possible	40
Section One Total	33

**2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY**

**2.1. Mode and rate of reproduction**

- A. No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or asexual reproduction). 0
- B. Limited reproduction (fewer than 10 viable seeds per plant AND no vegetative reproduction; if viability is not known, then maximum seed production is less than 100 seeds per plant and no vegetative reproduction) 1
- C. Moderate reproduction (fewer than 100 viable seeds per plant - if viability is not known, then maximum seed production is less than 1000 seeds per plant - OR limited successful vegetative spread documented) 2
- D. Abundant reproduction with vegetative asexual spread documented as one of the plants prime reproductive means OR more than 100 viable seeds per plant (if viability is not known, then maximum seed production reported to be greater than 1000 seeds per plant.) 4
- U. Unknown

Score 4

**Documentation:**

Describe key reproductive characteristics (including seeds per plant):  
Individual plants can produce over 1000 seeds/ year; can remain viable for at least five years.

Sources of information:

Swearingen & Adams, 2007

**2.2. Innate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair, buoyant fruits, pappus for wind-dispersal)**

- A. Does not occur (no long-distance dispersal mechanisms) 0
- B. Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of adaptations) 1
- C. Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant) 2
- D. Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent plant) 4
- U. Unknown

Score 2

**Documentation:**

Identify dispersal mechanisms:  
Small, flat seeds could easily be dispersed by numerous animals and flowing water.

Sources of information:

Moore, 2008.

**2.3. Potential to be spread by human activities (both directly and indirectly – possible mechanisms include: commercial sales, use as forage/revegetation, spread along highways, transport on boats, contaminated compost, land and vegetation)**

**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
Form originally created for use in New York. Indiana Form version date: November 1, 2010

---

management equipment such as mowers and excavators, etc.)

- A. Does not occur 0
- B. Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) 1
- C. Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) 2
- D. High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) 3
- U. Unknown

Score

**Documentation:**

Identify dispersal mechanisms:

Small, flat seeds could easily be dispersed by numerous animals and flowing water.

Sources of information:

Moore, 2008.

2.4. Characteristics that increase competitive advantage, such as shade tolerance, ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc.

- A. Possesses no characteristics that increase competitive advantage 0
- B. Possesses one characteristic that increases competitive advantage 3
- C. Possesses two or more characteristics that increase competitive advantage 6
- U. Unknown

Score

**Documentation:**

Evidence of competitive ability:

Shade tolerant, fast growth, low palatability to white-tailed deer

Sources of information:

Gleason & Cronquist, 1991; Brooklyn Botanic Garden, 2007; Swearingen & Adams, 2007; Eschtruth & Battles, 2008.

2.5. Growth vigor

- A. Does not form thickets or have a climbing or smothering growth habit 0
- B. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms 2
- U. Unknown

Score

**Documentation:**

Describe growth form:

Dominates herb layer; forms a thatch that persists beyond growing season.

Sources of information:

Swearingen & Adams, 2007

2.6. Germination/Regeneration

- A. Requires open soil or water and disturbance for seed germination, or regeneration from vegetative propagules. 0
- B. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions 2
- C. Can germinate/regenerate in existing vegetation in a wide range of conditions 3
- U. Unknown (No studies have been completed)

Score

**Documentation:**

**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
 Form originally created for use in New York. Indiana Form version date: November 1, 2010

Describe germination requirements:  
 Will germinate under numerous light and soil conditions.  
 Sources of information:  
 Swearingen & Adams, 2007.

2.7. Other species in the genus invasive in Indiana or elsewhere

- A. No 0
- B. Yes 3
- U. Unknown

Score 

0
---

Documentation:  
 Species:  
 Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008

Total Possible 

25
----

  
 Section Two Total 

20
----

**3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION**

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada (use same definition as Gleason & Cronquist which is: “The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario lying south of the 47th parallel of latitude”)

- A. No large stands (no areas greater than 1/4 acre or 1000 square meters) 0
- B. Large dense stands present in areas with numerous invasive species already present or disturbed landscapes 2
- C. Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas) 4
- U. Unknown

Score 

4
---

Documentation:  
 Identify reason for selection, or evidence of weedy history:  
 Plant is known and has been observed to invade forested areas where few other non-natives are noted.  
 Sources of information:  
 Swearingen & Adams, 2007; Moore, 2008.

3.2. Number of habitats the species may invade

- A. Not known to invade any natural habitats given at A2.2 0
- B. Known to occur in two or more of the habitats given at A2.2, with at least one a natural habitat. 1
- C. Known to occur in three or more of the habitats given at A2.2, with at least two a natural habitat. 2
- D. Known to occur in four or more of the habitats given at A2.2, with at least three a natural habitat. 4
- E. Known to occur in more than four of the habitats given at A2.2, with at least four a natural 6

**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
 Form originally created for use in New York. Indiana Form version date: November 1, 2010

habitat.  
 U. Unknown

Score

**Documentation:**

Identify type of habitats where it occurs and degree/type of impacts:  
 See A2.2.  
 Sources of information:

**3.3. Role of disturbance in establishment**

- A. Requires anthropogenic disturbances to establish. 0
- B. May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances. 2
- C. Can establish independent of any known natural or anthropogenic disturbances. 4
- U. Unknown

Score

**Documentation:**

Identify type of disturbance:  
 Mowing, tilling, foot traffic, and other soil disturbing activities as well as natural disturbances such as the scouring associated with flooding.  
 Sources of information:  
 Swearingen & Adams, 2007; Moore, 2008.

**3.4. Climate in native range**

- A. Native range does not include climates similar to Indiana 0
- B. Native range possibly includes climates similar to at least part of Indiana 1
- C. Native range includes climates similar to those in Indiana 3
- U. Unknown

Score

**Documentation:**

Describe what part of the native range is similar in climate to Indiana:  
 Europe.  
 Sources of information:  
 U.S.D.A., 2008; Brooklyn Botanic Garden, 2008

**3.5. Current introduced distribution in the northeastern USA and eastern Canada (see question 3.1 for definition of geographic scope )**

- A. Not known from the northeastern US and adjacent Canada 0
- B. Present as a non-native in one northeastern USA state and/or eastern Canadian province. 1
- C. Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces. 2
- D. Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 1 northeastern state or eastern Canadian province. 3
- E. Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 2 northeastern states or eastern Canadian provinces. 4
- U. Unknown

Score

**Documentation:**

Identify states and provinces invaded:  
 Connecticut, Delaware, District of Columbia, Kentucky, Illinois, Indiana, Maryland, Massachusetts, Missouri, New Jersey, New York, Ohio, Pennsylvania, Tennessee, Virginia,

**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
Form originally created for use in New York. Indiana Form version date: November 1, 2010

---

West Virginia  
Sources of information:  
See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  
Thieret, 2003; U.S.D.A., 2007.

3.6. Current introduced distribution of the species in natural areas in Indiana

- A. Present in no Indiana counties 0
- B. Present in 1-10 Indiana counties 1
- C. Present in 11-20 Indiana counties 2
- D. Present in 21-50 Indiana counties 3
- E. Present in more than 50 Indiana counties or on Federal noxious weed list 4
- U. Unknown

Score

Documentation:  
Describe distribution:  
See A1.1.  
Sources of information:

Total Possible   
Section Three Total

---

**4. DIFFICULTY OF CONTROL**

4.1. Seed banks

- A. Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make viable seeds or persistent propagules. 0
- B. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years 2
- C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years 3
- U. Unknown

Score

Documentation:  
Identify longevity of seed bank:  
At least five years  
Sources of information:  
Swearingen & Adams, 2007.

4.2. Vegetative regeneration

- A. No regrowth following removal of aboveground growth 0
- B. Regrowth from ground-level meristems 1
- C. Regrowth from extensive underground system 2
- D. Any plant part is a viable propagule 3
- U. Unknown

Score

Documentation:  
Describe vegetative response:  
Being a grass, the species possesses intercalary meristems and regrowth can therefore occur from above- and below ground tissues.



**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
Form originally created for use in New York. Indiana Form version date: November 1, 2010

Sources of information:  
Swearingen & Adams, 2007.

4.3. Level of effort required

- A. Management is not required: e.g., species does not persist without repeated anthropogenic disturbance. 0
- B. Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft<sup>2</sup>). 2
- C. Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). 3
- D. Management requires a major investment: e.g. more than 100 person-hours/year of manual effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. Eradication may be impossible (infestation as above). 4
- U. Unknown

Score 

4
---

Documentation:  
Identify types of control methods and time-term required:  
Hand removal with repeated follow up due to seeds that remain viable for many years  
Sources of information:  
Swearingen & Adams, 2007.

Total Possible 

10
----

  
Section Four Total 

7
---

**Total for 4 sections Possible**

100
-----

  
**Total for 4 sections**

84
----

**References for species assessment:**

Barden, L. 1987. Invasion of *Microstegium vimineum* (Poaceae), an exotic, annual, shade-tolerant, C-4 grass, into a North Carolina floodplain. *Amer. Midl. Naturalist* 118 (1): 40-45.

Barden, L. 1991. Element Stewardship Abstract: *Microstegium vimineum*. The Nature Conservancy. Arlington, Virginia, United States. 6 p.

Brooklyn Botanic Garden. 2008. AILANTHUS database. [Accessed on 8 February 2008].

Claridge, K. & S. B. Franklin. 2003. Compensation and plasticity in an invasive plant species. *Biol. Invas.* 4: 339-347.

Cole, P. G. & J. F. Weltzin. 2004. Environmental correlates of the distribution and abundance of *Microstegium vimineum*, in east Tennessee. *S.E. Naturalist (Steuben)* 3: 545-562.

Cole, P. G. & J. F. Weltzin. 2005. Light limitation creates patchy distribution of an invasive grass in eastern deciduous forests. *Biol. Invas.* 7: 477-488.

Ehrenfeld, J. G., P. Kourtev & W. Huang. 2001. Changes in soil functions following invasions of exotic understory plants in deciduous forests. *Ecol. Appl.* 11: 1287-1300.

# INDIANA

## NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE

Form originally created for use in New York. Indiana Form version date: November 1, 2010

---

Eschtruth, A.K. and J.J. Battles. 2008. Acceleration of exotic plant invasion in a forested ecosystem by a generalist herbivore. *Conserv. Biol.* In Press

Flory, S. L. 2011. Invasive grass (*Microstegium vimineum*) increases fire intensity and reduced tree regeneration in eastern forests. Presentation at the 2011 Natural Areas Association Conference.

Horton, J. L.D. Mabe (original application development), Florida Center for Community Design and Research. University of South Florida]. & H. S. Neufeld. 1998. Photosynthetic responses of *Microstegium vimineum* (Trin.) A. Camus, a shade-tolerant, C4 grass, to variable light environments. *Oecologia* 114: 11-19.

Hunt, D. M. & R. E. Zaremba. 1992. The northeastward spread of *Microstegium vimineum* (Poaceae) into New York and adjacent states. *Rhodora* 94: 167-170.

Kourtev, P. S., J. G. Ehrenfeld, & M. Häggblom. 2002. Exotic plant species alter the microbial community structure and function in the soil. *Ecology* 83(11): 3152-3166.

LaFleur, A. 1996. Invasive plant information sheet: Japanese stiltgrass. The Nature Conservancy, Connecticut Chapter. Hartford, Connecticut, United States.

Moore, G. 2008. Invasiveness ranking system for non-native plants of New York: *Microstegium vimineum*. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY.

Redman, D. E. 1995. Distribution and habitat types for Nepal microstegium [*Microstegium vimineum* (Trin.) Camus] in Maryland and the District of Columbia. *Castanea* 60(3): 270-275.

Swearingen, J. M. & S. Adams. 2007. Fact Sheet: Japanese stiltgrass [online]. Plant Conservation Alliance. <<http://www.nps.gov/plants/ALIEN/fact/pdf/mivi1.pdf>> [Accessed 15 October 2007].

United States Department of Agriculture, National Resources Conservation Service. 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, Louisiana [Accessed on 31 December 2007].

Winter, K., M. R. Schmitt & G. E. Edwards. 1982. *Microstegium vimineum*, a shade adapted C-4 grass. *Pl. Sci. Lett.* 24: 311-318.

Weldy, T. and D. Werier. 2005. New York Flora Atlas. [S.M. Landry, K.N. Campbell, and L.D. Mabe (original application development), Florida Center for Community Design and Research. University of South Florida]. New York Flora Association, Albany, New York. [Accessed on 31 December 2007].

**Citation:** This IN ranking form may be cited as: Jacquart, E.M. 2012. Invasiveness ranking system for non-native plants of Indiana. Unpublished. Invasive Plant Advisory Committee (IPAC) to the Indiana Invasive Species Council, Indianapolis, IN.

**Acknowledgments:** The IN ranking form is an adaptation for Indiana use of the form created for New York by Jordan et al. (2009), cited below. Documentation for species assessed for New York are used for Indiana where they are applicable. The Invasive Plant Advisory Committee was created by the Indiana Invasive Species Council in October 2010, and is made up of the original members of the Indiana Invasive Plant Assessment Working Group (IPSAWG). Original members of IPSAWG included representatives of the The Nature Conservancy; Indiana Native Plant and Wildflower Society; Indiana Nursery and Landscape Association; Indiana Chapter of the American Society of Landscape Architects; Indiana Forage Council; Indiana Wildlife Federation; Indiana State Beekeepers Association; Indiana Beekeeper's Association; Department of Natural Resources; Hoosier National Forest; Indiana Academy of Science; Natural Resources Conservation Service; Indiana Department of Environmental Management;

**INDIANA**  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE  
Form originally created for use in New York. Indiana Form version date: November 1, 2010

---

Indiana Department of Transportation; Purdue Cooperative Extension Service; Seed Administrator, Office of the Indiana State Chemist.

**References for the Indiana ranking form:**

Jordan, M.J., G. Moore, and T.W. Weldy. 2009. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY.

**References for the New York ranking form:**

Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: [http://akweeds.uaa.alaska.edu/akweeds\\_ranking\\_page.htm](http://akweeds.uaa.alaska.edu/akweeds_ranking_page.htm).

Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).

Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. <http://www.natureserve.org/getData/plantData.jsp>

Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. *Invasive Plant Science and Management* 1:36–49

Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M. Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at [www.caleppc.org](http://www.caleppc.org) and [www.swvma.org](http://www.swvma.org). California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.

Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. *Science for Conservation* 209. New Zealand Department of Conservation. 1-23 pp.