

INDIANA

NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
 Form originally created for use in New York
 Indiana Form version date: November 1, 2010

Scientific name: Carduus acanthoides USDA Plants Code: CAAC
 Common names: Spiny Plumeless Thistle, Plumeless Thistle, Weltd Thistle
 Native distribution: Southern Europe and western Asia
 Date assessed: 7/2/2013
 Assessors: Zach Deitch, Ellen Jacquart
 Reviewers: Scott Namestnik, Stuart Orr
 Date Approved: 8/8/2013

Indiana Invasiveness Rank:

Invasiveness Ranking Summary (see details under appropriate sub-section)		Total (Total Answered*) Possible	Total
1	Ecological impact	40 (<u>20</u>)	14
2	Biological characteristic and dispersal ability	25 (<u>19</u>)	18
3	Ecological amplitude and distribution	25 (<u>25</u>)	20
4	Difficulty of control	10 (<u>10</u>)	7
	Outcome score	100 (<u>74</u>) ^b	59 ^a
	Relative maximum score [†]		79.7
	Indiana Invasiveness Rank [§]	High	

* For questions answered "unknown" do not include point value in "Total Answered Points Possible." If "Total Answered Points Possible" is less than 70.00 points, then the overall invasive rank should be listed as "Unknown."
[†]Calculated as 100(a/b) to two decimal places.
[§]Very High >80.00; High 70.00–80.00; Moderate 50.00–69.99; Low 40.00–49.99; Insignificant <40.00

A. DISTRIBUTION (KNOWN/POTENTIAL):

A1 Has this species been documented to persist without cultivation in IN? (reliable source; voucher not required)


Yes – continue to A2.2

No – continue to A2.1

A2 What is the likelihood that this species will occur and persist outside of cultivation given the climate in Indiana? (obtain from occurrence data in other states with similar climates)

Likely – continue to A3

Not likely – stop here. There is no need to assess the species



INDIANA

NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
 Form originally created for use in New York
 Indiana Form version date: November 1, 2010

Documentation:

Sources of information: Range maps compiled from PLANTS database, <http://plants.usda.gov/java/>; Indiana CAPS database, <http://extension.entm.purdue.edu/CAPS/index.html>; Indiana IPSAWG reports (unpublished); and EDDMapS reports, <http://eddmaps.org/>

A3 Describe the potential or known suitable habitats within Indiana (underlined). Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

Aquatic Habitats Rivers/streams Natural lakes and ponds Reservoirs/impoundments*	Wetland Habitats Marshes Fens Bogs Shrub swamps Forested wetlands/riparian <u>Beaches/dunes</u> <u>Ditches*</u>	Upland Habitats Forest <u>Savannas</u> <u>Barrens</u> <u>Prairies</u> Cultivated* <u>Old Fields*</u> <u>Roadsides*</u>
---	--	---

Other potential or known suitable habitats within Indiana: glade communities, buffer zones, restorations, abandoned agricultural land, dumps, fencerows, pastures, canopy gaps and open spaces in high quality natural areas.

Documentation: *roadsides, disturbed areas, ditch banks, and old fields and spreads into high quality prairies, meadows, and pastures.*

Sources of information:
 WDNR-Invasive Species.
 Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

B. INVASIVENESS RANKING

Questions apply to areas similar in climate and habitats to Indiana unless specified otherwise.

1. ECOLOGICAL IMPACT

1.1. Impact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire regime, geomorphological changes (erosion, sedimentation rates), hydrologic regime, nutrient and mineral dynamics, light availability, salinity, pH)

- A. No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the northeast for >100 years. 0
- B. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability) 3
- C. Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl) 7
- D. Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species) 10
- U. Unknown

Score U

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Documentation:

Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)

Infests low quality areas first such as roadsides, disturbed areas, ditch banks, and old fields and spreads into high quality prairies. When in meadows and pastures, grazing animals avoid plumeless thistle and focus on native plants giving the invasive the upper hand. Prairie and grassland communities provide ecosystem services (carbon sequestration).

Fire will not push through heavy infestations. May distract pollinators from native species.

Sources of information:

WDNR-Invasive Species.

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

1.2. Impact on Natural Community Structure

- A. No perceived impact; establishes in an existing layer without influencing its structure 0
- B. Influences structure in one layer (e.g., changes the density of one layer) 3
- C. Significant impact in at least one layer (e.g., creation of a new layer or elimination of an existing layer) 7
- D. Major alteration of structure (e.g., covers canopy, eradicating most or all layers below) 10
- U. Unknown

Score

Documentation:

Identify type of impact or alteration:

Can form monotypic vegetation stands. However, this does not affect structure so much as it affects composition.

Sources of information:

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

1.3. Impact on Natural Community Composition

- A. No perceived impact; causes no apparent change in native populations 0
- B. Influences community composition (e.g., reduces the number of individuals in one or more native species in the community) 3
- C. Significantly alters community composition (e.g., produces a significant reduction in the population size of one or more native species in the community) 7
- D. Causes major alteration in community composition (e.g., results in the extirpation of one or several native species, reducing biodiversity or change the community composition towards species exotic to the natural community) 10
- U. Unknown

Score

Documentation:

Identify type of impact or alteration:

When in meadows and pastures, grazing animals avoid plumeless thistle and focus on native plants giving the invasive the upper hand.

Reduces native species density and diversity.

Can form monotypic stands.

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Sources of information:
WDNR-Invasive Species.
Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

1.4. Impact on other species or species groups (cumulative impact of this species on the animals, fungi, microbes, and other organisms in the community it invades. Examples include reduction in nesting/foraging sites; reduction in habitat connectivity; injurious components such as spines, thorns, burrs, toxins; suppresses soil/sediment microflora; interferes with native pollinators and/or pollination of a native species; hybridizes with a native species; hosts a non-native disease which impacts a native species)

- | | | |
|----|--|----|
| A. | Negligible perceived impact | 0 |
| B. | Minor impact | 3 |
| C. | Moderate impact | 7 |
| D. | Severe impact on other species or species groups | 10 |
| U. | Unknown | |

Score 7

Documentation:
Identify type of impact or alteration:
When in meadows and pastures, grazing animals avoid plumeless thistle and focus on native plants giving the invasive the upper hand.

Prairie and grassland communities provide ecosystem services (carbon sequestration) and habitat for arthropods and birds, and other wildlife.
Sources of information:
WDNR-Invasive Species.
Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

Total Possible	20
Section One Total	14

2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY

2.1. Mode and rate of reproduction

- | | | |
|----|---|---|
| A. | No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or asexual reproduction). | 0 |
| B. | Limited reproduction (fewer than 10 viable seeds per plant AND no vegetative reproduction; if viability is not known, then maximum seed production is less than 100 seeds per plant and no vegetative reproduction) | 1 |
| C. | Moderate reproduction (fewer than 100 viable seeds per plant - if viability is not known, then maximum seed production is less than 1000 seeds per plant - OR limited successful vegetative spread documented) | 2 |
| D. | Abundant reproduction with vegetative asexual spread documented as one of the plants prime reproductive means OR more than 100 viable seeds per plant (if viability is not known, then maximum seed production reported to be greater than 1000 seeds per plant.) | 4 |
| U. | Unknown | |

Score 4

Documentation:
Describe key reproductive characteristics (including seeds per plant):
Up to 10,000 seeds per plant.

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Biennial. Most common in well-drained or sandy, fertile soils with neutral or slightly acidic pH. Seeds are dispersed 1 - 3 weeks after flowering.

Sources of information:

WDNR-Invasive Species.

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

2.2. Innate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair, buoyant fruits, pappus for wind-dispersal)

- A. Does not occur (no long-distance dispersal mechanisms) 0
- B. Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of adaptations) 1
- C. Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant) 2
- D. Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent plant) 4
- U. Unknown

Score

Documentation:

Identify dispersal mechanisms:

Wind, humans, and vehicles are the main way the seeds get dispersed.

Sources of information:

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

2.3. Potential to be spread by human activities (both directly and indirectly – possible mechanisms include: commercial sales, use as forage/revegetation, spread along highways, transport on boats, contaminated compost, land and vegetation management equipment such as mowers and excavators, etc.)

- A. Does not occur 0
- B. Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) 1
- C. Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) 2
- D. High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) 3
- U. Unknown

Score

Documentation:

Identify dispersal mechanisms:

Intentional:

Unintentional: Bird Animal Vehicles/Human

Wind Water Other: Seeds can adhere to farm machinery, vehicles, agricultural products like hay and straw.

Sources of information:

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

2.4. Characteristics that increase competitive advantage, such as shade tolerance, ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc.

- A. Possesses no characteristics that increase competitive advantage 0
- B. Possesses one characteristic that increases competitive advantage 3

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

- C. Possesses two or more characteristics that increase competitive advantage 6
U. Unknown

Score

Documentation:

Evidence of competitive ability:

Most common in well-drained or sandy, fertile soils with neutral or slightly acidic pH. It is a prolific seeder and seeds are dispersed 1 - 3 weeks after flowering.

Sources of information:

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

2.5. Growth vigor

- A. Does not form thickets or have a climbing or smothering growth habit 0
B. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms 2
U. Unknown

Score

Documentation:

Describe growth form: *Can form weedy monotypic stands that crowd out native plants.*

Sources of information:

Kimmel, 2012.

2.6. Germination/Regeneration

- A. Requires open soil or water and disturbance for seed germination, or regeneration from vegetative propagules. 0
B. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions 2
C. Can germinate/regenerate in existing vegetation in a wide range of conditions 3
U. Unknown (No studies have been completed)

Score

Documentation:

Describe germination requirements:

Dry, open or partially shaded areas. Most common in well-drained or sandy, fertile soils with neutral or slightly acidic pH. Flowers May to July. Seeds are dispersed 1 - 3 weeks after flowering.

Sources of information:

Plumeless Thistle (*Carduus acanthoides*).

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

2.7. Other species in the genus invasive in Indiana or elsewhere

- A. No 0
B. Yes 3
U. Unknown

Score

Documentation:

Carduus nutans is invasive in Indiana.

Species:

Total Possible	19
Section Two Total	18

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada (use same definition as Gleason & Cronquist which is: “The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario lying south of the 47th parallel of latitude”)

- A. No large stands (no areas greater than 1/4 acre or 1000 square meters) 0
- B. Large dense stands present in areas with numerous invasive species already present or disturbed landscapes 2
- C. Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas) 4
- U. Unknown

Score

4

Documentation:

Identify reason for selection, or evidence of weedy history:

Can form weedy monotypic stands that crowd out native plants and can resist fire if large enough.

Infests low quality areas first such as roadsides, disturbed areas, ditch banks, and old fields and spreads into high quality prairies. When in meadows and pastures, grazing animals avoid plumeless thistle and focus on native plants giving the invasive the upper hand.

Sources of information:

Kimmel, 2012.

WDNR-Invasive Species.

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*

3.2. Number of habitats the species may invade

- A. Not known to invade any natural habitats given at A2.2 0
- B. Known to occur in two or more of the habitats given at A2.2, with at least one a natural habitat. 1
- C. Known to occur in three or more of the habitats given at A2.2, with at least two a natural habitat. 2
- D. Known to occur in four or more of the habitats given at A2.2, with at least three a natural habitat. 4
- E. Known to occur in more than four of the habitats given at A2.2, with at least four a natural habitat. 6
- U. Unknown

Score

6

Documentation:

Identify type of habitats where it occurs and degree/type of impacts:

Seven habitats with four being natural identified in A3.

Sources of information:

See A3.

3.3. Role of disturbance in establishment

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

- A. Requires anthropogenic disturbances to establish. 0
- B. May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances. 2
- C. Can establish independent of any known natural or anthropogenic disturbances. 4
- U. Unknown

Score

Documentation:

Identify type of disturbance:

Dry, open or partially shaded areas with disturbance.

Infests low quality areas first such as roadsides, disturbed areas, ditch banks, and old fields and spreads into high quality prairies. When in meadows and pastures, grazing animals avoid plumeless thistle and focus on native plants giving the invasive the upper hand.

Sources of information:

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

3.4. Climate in native range

- A. Native range does not include climates similar to Indiana 0
- B. Native range possibly includes climates similar to at least part of Indiana 1
- C. Native range includes climates similar to those in Indiana 3
- U. Unknown

Score

Documentation:

Describe what part of the native range is similar in climate to Indiana:

Invasive throughout the United States expanding into open areas. Most common in well-drained or sandy, fertile soils with neutral or slightly acidic pH. Requires cold period to induce reproductive stage.

Sources of information:

Wisconsin Invasive Plant Assessment- *Carduus acanthoides*.

USDA, NRCS. 2007.

3.5. Current introduced distribution in the northeastern USA and eastern Canada (see question 3.1 for definition of geographic scope)

- A. Not known from the northeastern US and adjacent Canada 0
- B. Present as a non-native in one northeastern USA state and/or eastern Canadian province. 1
- C. Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces. 2
- D. Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 1 northeastern state or eastern Canadian province. 3
- E. Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 2 northeastern states or eastern Canadian provinces. 4
- U. Unknown

Score

Documentation:

Identify states and provinces invaded:

Found in British Columbia, Ontario, Alberta, Quebec and Nova Scotia. Widespread in North America.

Sources of information:

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Carduus genus.
Kimmel, 2012.

3.6. Current introduced distribution of the species in natural areas in Indiana

- | | |
|---|---|
| A. Present in no Indiana counties | 0 |
| B. Present in 1-10 Indiana counties | 1 |
| C. Present in 11-20 Indiana counties | 2 |
| D. Present in 21-50 Indiana counties | 3 |
| E. Present in more than 50 Indiana counties or on Federal noxious weed list | 4 |
| U. Unknown | |

Score

Documentation:
Describe distribution:
Documented in 0 counties of Indiana.
Sources of information:
See A1

Total Possible	25
Section Three Total	20

4. DIFFICULTY OF CONTROL

4.1. Seed banks

- | | |
|---|---|
| A. Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make viable seeds or persistent propagules. | 0 |
| B. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years | 2 |
| C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years | 3 |
| U. Unknown | |

Score

Documentation:
Seed can remain viable in the soil for over 10 years.
Sources of information:
WDNR- Invasive Species.

4.2. Vegetative regeneration

- | | |
|--|---|
| A. No regrowth following removal of aboveground growth | 0 |
| B. Regrowth from ground-level meristems | 1 |
| C. Regrowth from extensive underground system | 2 |
| D. Any plant part is a viable propagule | 3 |
| U. Unknown | |

Score

Documentation:
Describe vegetative response:
Taproot. Biennial with basal rosette in first growing stage.
Sources of information:
WDNR- Invasive Species.

4.3. Level of effort required

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

- A. Management is not required: e.g., species does not persist without repeated anthropogenic disturbance. 0
- B. Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft²). 2
- C. Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). 3
- D. Management requires a major investment: e.g. more than 100 person-hours/year of manual effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. Eradication may be impossible (infestation as above). 4
- U. Unknown

Score

3

Documentation:
Identify types of control methods and time-term required:

Can be eradicated in several growing seasons.

Mechanical: Close mowing of second year plants twice per growing season just before flowering. Cut or mow at early bud both times. Cut plants with sharp shovel 1-2" below soil surface before flowering.

Chemical: Spot spray rosettes in fall with 2, 4-D ester; foliar spray with clopyralid or metsulfuron methyl.

Sources of information:
WDNR- Invasive Species.

Total Possible

10

Section Four Total

7

Total for 4 sections Possible

100

Total for 4 sections

89

References for species assessment:

“Carduus genus”. *California Department of Food Agriculture*.
<http://www.cdfa.ca.gov/plant/ipc/weedinfo/carduus.htm>

Kimmel, N. 2012. “Plumeless Thistle (Carduus acanthoides)”. *Alberta Weed Monitoring Network*.
[http://www1.agric.gov.ab.ca/\\$department/deptdocs.nsf/all/prm13967](http://www1.agric.gov.ab.ca/$department/deptdocs.nsf/all/prm13967)

Plumeless Thistle (Carduus acanthoides). “ODA Plant Programs, Noxious Weed Control”.
http://www.oregon.gov/ODA/PLANT/weeds/Pages/profile_plumelessthistle.aspx

USDA, NRCS. 2007. The PLANTS Database (<http://plants.usda.gov>, 16 March 2007). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

INDIANA

NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE

Form originally created for use in New York

Indiana Form version date: November 1, 2010

WDNR - Invasive Species. <http://dnr.wi.gov/topic/Invasives/fact/PlumelessThistle.html>

Wisconsin Invasive Plant Assessment for *Carduus acanthoides*.

http://dnr.wi.gov/topic/Invasives/documents/classification/LR_Carduus_acanthoides.pdf Date Accessed: 19 June 2013.

Wisconsin State Herbarium. 2007. WISFLORA: Wisconsin Vascular Plant Species

(<http://www.botany.wisc.edu/wisflora/>). Dept. Botany, Univ. Wisconsin, Madison, WI 53706-1381 USA.

Citation: This IN ranking form may be cited as: Jacquart, E.M. 2011. Invasiveness ranking system for non-native plants of Indiana. Unpublished. Invasive Plant Advisory Committee (IPAC) to the Indiana Invasive Species Council, Indianapolis, IN.

Acknowledgments: The IN ranking form is an adaptation for Indiana use of the form created for New York by Jordan et al. (2009), cited below. Documentation for species assessed for New York are used for Indiana where they are applicable. The Invasive Plant Advisory Committee was created by the Indiana Invasive Species Council in October 2010, and is made up of the original members of the Indiana Invasive Plant Assessment Working Group (IPSAWG). Original members of IPSAWG included representatives of the The Nature Conservancy; Indiana Native Plant and Wildflower Society; Indiana Nursery and Landscape Association; Indiana Chapter of the American Society of Landscape Architects; Indiana Forage Council; Indiana Wildlife Federation; Indiana State Beekeepers Association; Indiana Beekeeper's Association; Department of Natural Resources; Hoosier National Forest; Indiana Academy of Science; Natural Resources Conservation Service; Indiana Department of Environmental Management; Indiana Department of Transportation; Purdue Cooperative Extension Service; Seed Administrator, Office of the Indiana State Chemist.

References for the Indiana ranking form:

Jordan, M.J., G. Moore, and T.W. Weldy. 2009. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY.

References for the New York ranking form:

Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: http://akweeds.uaa.alaska.edu/akweeds_ranking_page.htm.

Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).

Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. <http://www.natureserve.org/getData/plantData.jsp>

Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. *Invasive Plant Science and Management* 1:36-49

Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M. Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE

Form originally created for use in New York

Indiana Form version date: November 1, 2010

www.caleppc.org and www.swvma.org. California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.

Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. Science for Conservation 209. New Zealand Department of Conservation. 1-23 pp.