

INDIANA

NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Scientific name: Dipsacus fullonum L. USDA Plants Code: DIFU2
 Common names: common teasel
 Native distribution: Europe, temperate Asia, northern Africa
 Date assessed: August 20, 2012
 Assessors: Ellen Jacquart
 Reviewers: Stuart Orr, Brenda Howard, Ken Collins
 Date Approved: September 21, 2012

Indiana Invasiveness Rank: High (Relative Maximum Score 70.00-80.00)

Invasiveness Ranking Summary (see details under appropriate sub-section)		Total (Total Answered*) Possible	Total
1	Ecological impact	40 (30)	17
2	Biological characteristic and dispersal ability	25 (25)	19
3	Ecological amplitude and distribution	25 (25)	23
4	Difficulty of control	10 (10)	7
	Outcome score	100 (90) ^b	66 ^a
	Relative maximum score [†]		73.33
	Indiana Invasiveness Rank [§]	High (Relative Maximum Score 70.00-80.00)	

* For questions answered “unknown” do not include point value in “Total Answered Points Possible.” If “Total Answered Points Possible” is less than 70.00 points, then the overall invasive rank should be listed as “Unknown.”

[†]Calculated as 100(a/b) to two decimal places.

[§]Very High >80.00; High 70.00–80.00; Moderate 50.00–69.99; Low 40.00–49.99; Insignificant <40.00


A. DISTRIBUTION (KNOWN/POTENTIAL):

A1 Has this species been documented to persist without cultivation in IN? (reliable source; voucher not required)

<input checked="" type="checkbox"/>	Yes – continue to A2.2
<input type="checkbox"/>	No – continue to A2.1

A2 What is the likelihood that this species will occur and persist outside of cultivation given the climate in Indiana? (obtain from occurrence data in other states with similar climates)

<input checked="" type="checkbox"/>	Likely – continue to A3
<input type="checkbox"/>	Not likely – stop here. There is no need to assess the species



INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Documentation:

Sources of information: Range maps compiled from PLANTS database, <http://plants.usda.gov/java/>; Indiana CAPS database, <http://extension.entm.purdue.edu/CAPS/index.html>; Indiana IPSAWG reports (unpublished); and EDDMapS reports, <http://eddmaps.org/>

A3 Describe the potential or known suitable habitats within Indiana (underlined). Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

Aquatic Habitats	Wetland Habitats	Upland Habitats
Rivers/streams	Marshes	Forest
Natural lakes and ponds	Fens	<u>Savannas</u>
<u>Reservoirs/impoundments*</u>	Bogs	<u>Barrens</u>
	Shrub swamps	<u>Prairies</u>
	Forested wetlands/riparian	Cultivated*
	Beaches/dunes	<u>Old Fields*</u>
	Ditches*	<u>Roadsides*</u>

Other potential or known suitable habitats within Indiana:
No additional habitats.

Documentation:

Sources of information:
Weber, 2003; Snyder & Kaufman, 2004; Fellows & Grauver, 2006; Gucker, 2009, Jacquart personal observation, Orr personal observation.

B. INVASIVENESS RANKING

Questions apply to areas similar in climate and habitats to Indiana unless specified otherwise.

1. ECOLOGICAL IMPACT

1.1. Impact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire regime, geomorphological changes (erosion, sedimentation rates), hydrologic regime, nutrient and mineral dynamics, light availability, salinity, pH)

- A. No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the northeast for >100 years. 0
- B. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability) 3
- C. Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl) 7
- D. Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species) 10
- U. Unknown

Score

U

Documentation:

Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)
Like cut-leaf teasel, there are no reports of significant impact on ecosystem processes or system wide parameters.

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Sources of information:
Fellows & Grauver, 2006.

1.2. Impact on Natural Community Structure

- | | | |
|----|--|----|
| A. | No perceived impact; establishes in an existing layer without influencing its structure | 0 |
| B. | Influences structure in one layer (e.g., changes the density of one layer) | 3 |
| C. | Significant impact in at least one layer (e.g., creation of a new layer or elimination of an existing layer) | 7 |
| D. | Major alteration of structure (e.g., covers canopy, eradicating most or all layers below) | 10 |
| U. | Unknown | |

Score 7

Documentation:

Identify type of impact or alteration:
Dense leaves can shade out other vegetation (Werner, 1972; Fellows & Grauver, 2006).
Sources of information:
Werner, 1972; Fellows & Grauver, 2006.

1.3. Impact on Natural Community Composition

- | | | |
|----|---|----|
| A. | No perceived impact; causes no apparent change in native populations | 0 |
| B. | Influences community composition (e.g., reduces the number of individuals in one or more native species in the community) | 3 |
| C. | Significantly alters community composition (e.g., produces a significant reduction in the population size of one or more native species in the community) | 7 |
| D. | Causes major alteration in community composition (e.g., results in the extirpation of one or several native species, reducing biodiversity or change the community composition towards species exotic to the natural community) | 10 |
| U. | Unknown | |

Score 7

Documentation:

Identify type of impact or alteration:
Smaller stands simply reduce the number of native individuals for a species in an area, whereas larger, dense stands can significantly reduce numbers of plant species. Also, in New Jersey, the species has been reported to have invaded limestone fens and caused the reduction or extirpation of several rare plant species, including spreading globe flower (*Trollius laxus subsp. laxus*) and sessile water speedwell (*Veronica catenata*) (Snyder & Kaufman, 2004).
Sources of information:
Snyder & Kaufman, 2004.

1.4. Impact on other species or species groups (cumulative impact of this species on the animals, fungi, microbes, and other organisms in the community it invades.

Examples include reduction in nesting/foraging sites; reduction in habitat connectivity; injurious components such as spines, thorns, burrs, toxins; suppresses soil/sediment microflora; interferes with native pollinators and/or pollination of a native species; hybridizes with a native species; hosts a non-native disease which impacts a native species)

- | | | |
|----|--|----|
| A. | Negligible perceived impact | 0 |
| B. | Minor impact | 3 |
| C. | Moderate impact | 7 |
| D. | Severe impact on other species or species groups | 10 |
| U. | Unknown | |

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Score

3

Documentation:

Identify type of impact or alteration:

Plant has spines on leaves to discourage grazing by large herbivores (Werner, 1975). Teasel seeds were consumed by game birds such as Californian quail and ring-necked pheasants; and rodents (Knight et al., 1979; Mittelbach & Gross, 1984). One experiment suggest that invertebrates caught in its water filled leaf bases could increase the plant seed set and increase seed mass:biomass ratio (Shaw & Shackleton, 2011).

Sources of information:

Werner, 1975; Knight et al. 1979; Mittelbach & Gross, 1984; Shaw & Shackleton, 2011.

Total Possible

30

Section One Total

17

2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY

2.1. Mode and rate of reproduction

- A. No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or asexual reproduction). 0
- B. Limited reproduction (fewer than 10 viable seeds per plant AND no vegetative reproduction; if viability is not known, then maximum seed production is less than 100 seeds per plant and no vegetative reproduction) 1
- C. Moderate reproduction (fewer than 100 viable seeds per plant - if viability is not known, then maximum seed production is less than 1000 seeds per plant - OR limited successful vegetative spread documented) 2
- D. Abundant reproduction with vegetative asexual spread documented as one of the plants prime reproductive means OR more than 100 viable seeds per plant (if viability is not known, then maximum seed production reported to be greater than 1000 seeds per plant.) 4
- U. Unknown

Score

4

Documentation:

Describe key reproductive characteristics (including seeds per plant):

Reproduces entirely by seed but it can regenerate after sustaining damage (Gucker, 2009). Werner calculated seed production from her data as: 854.6 ± 375.7 seeds per inflorescence; and finding 3.9 ± 2.4 inflorescences per plant from a roadside population ($n = 15$) from Michigan and then multiplied 0.80 for % seed viability, but it can be much higher than 3000 (commonly cited number) per plant; seed viability is very high when allowed to cross-pollinate and much lower if self-pollinated (Werner, 1975).

Sources of information:

Werner, 1975; Gucker, 2009.

2.2. Innate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair, buoyant fruits, pappus for wind-dispersal)

- A. Does not occur (no long-distance dispersal mechanisms) 0
- B. Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of adaptations) 1
- C. Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant) 2
- D. Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent plant) 4
- U. Unknown

Score

1

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Documentation:

Identify dispersal mechanisms:

Werner cites, from a previous study, that 99.9% of the seeds fell passively and deposited themselves within 1.5 m of the parent plant. Occasional long distance dispersal by water or by people may occur (Werner, 1975; Fellows & Grauver, 2006; Gucker, 2009).

Sources of information:

Werner, 1975; Weber, 2003; Fellows & Grauver, 2006; Gucker, 2009.

2.3. Potential to be spread by human activities (both directly and indirectly – possible mechanisms include: commercial sales, use as forage/revegetation, spread along highways, transport on boats, contaminated compost, land and vegetation management equipment such as mowers and excavators, etc.)

- A. Does not occur 0
- B. Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) 1
- C. Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) 2
- D. High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) 3
- U. Unknown

Score

Documentation:

Identify dispersal mechanisms:

Readily dispersed by mowing equipment; also occasionally sold for cultivation and in dried flower displays.

Sources of information:

Donaldson & Rafferty, 2002; Snyder & Kaufman, 2004; Fellows & Grauver, 2006; Gucker, 2009

2.4. Characteristics that increase competitive advantage, such as shade tolerance, ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc.

- A. Possesses no characteristics that increase competitive advantage 0
- B. Possesses one characteristic that increases competitive advantage 3
- C. Possesses two or more characteristics that increase competitive advantage 6
- U. Unknown

Score

Documentation:

Evidence of competitive ability:

Perennial or biennial monocarp (i.e., dies after setting seed); deep taproots may extend below the roots of grasses found in abandoned fields (Werner, 1972)

Sources of information:

Werner, 1972; Fellows & Grauver, 2006.

2.5. Growth vigor

- A. Does not form thickets or have a climbing or smothering growth habit 0
- B. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms 2
- U. Unknown

Score

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Documentation:

Describe growth form:

Forms a very dense tall thickety layer above shorter vegetation.

Sources of information:

Snyder & Kaufman, 2004; Grauver, 2006; Jacquart pers. obs.

2.6. Germination/Regeneration

- A. Requires open soil or water and disturbance for seed germination, or regeneration from vegetative propagules. 0
- B. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions 2
- C. Can germinate/regenerate in existing vegetation in a wide range of conditions 3
- U. Unknown (No studies have been completed)

Score

Documentation:

Describe germination requirements:

Germinates in existing vegetation in a wide range of conditions, but seedling survival is best with moderate amounts of litter or beneath sparse vegetation (Hubbell & Werner, 1979). Uva, cited by Gucker, frequently found common teasel in damp and rich soils.

Sources of information:

Hubbell & Werner, 1979; & Uva et al., 1997; Grauver, 2006; Gucker, 2009; author's pers. obs.

2.7. Other species in the genus invasive in Indiana or elsewhere

- A. No 0
- B. Yes 3
- U. Unknown

Score

Documentation:

Species:

Dipsacus laciniatus considered invasive in Indiana

Total Possible	25
Section Two Total	19

3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada (use same definition as Gleason & Cronquist which is: "The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario lying south of the 47th parallel of latitude")

- A. No large stands (no areas greater than 1/4 acre or 1000 square meters) 0
- B. Large dense stands present in areas with numerous invasive species already present or disturbed landscapes 2
- C. Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas) 4
- U. Unknown

Score

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Documentation:

Identify reason for selection, or evidence of weedy history:

Large stands can occur over 0.25 acres sometimes in areas lacking other invasives.

Sources of information:

Snyder & Kaufman, 2004; Gravuer, 2006, Jacquart personal observation.

3.2. Number of habitats the species may invade

- | | | |
|----|---|---|
| A. | Not known to invade any natural habitats given at A2.2 | 0 |
| B. | Known to occur in two or more of the habitats given at A2.2, with at least one a natural habitat. | 1 |
| C. | Known to occur in three or more of the habitats given at A2.2, with at least two a natural habitat. | 2 |
| D. | Known to occur in four or more of the habitats given at A2.2, with at least three a natural habitat. | 4 |
| E. | Known to occur in more than four of the habitats given at A2.2, with at least four a natural habitat. | 6 |
| U. | Unknown | |

Score

Documentation:

Identify type of habitats where it occurs and degree/type of impacts:

See A2.2.

Sources of information:

Snyder & Kaufman, 2004; Grauver, 2006; Brooklyn Botanic Garden, 2009

3.3. Role of disturbance in establishment

- | | | |
|----|--|---|
| A. | Requires anthropogenic disturbances to establish. | 0 |
| B. | May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances. | 2 |
| C. | Can establish independent of any known natural or anthropogenic disturbances. | 4 |
| U. | Unknown | |

Score

Documentation:

Identify type of disturbance:

Usually found in disturbed areas, but also reported from undisturbed areas such as prairies.

Sources of information:

Snyder & Kaufman, 2005; Grauver, 2006; Jacquart personal observation

3.4. Climate in native range

- | | | |
|----|---|---|
| A. | Native range does not include climates similar to Indiana | 0 |
| B. | Native range possibly includes climates similar to at least part of Indiana | 1 |
| C. | Native range includes climates similar to those in Indiana | 3 |
| U. | Unknown | |

Score

Documentation:

Describe what part of the native range is similar in climate to Indiana:

Temperate Europe and Asia.

Sources of information:

Grauver, 2006; Brooklyn Botanic Garden, 2009.

3.5. Current introduced distribution in the northeastern USA and eastern Canada (see

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

question 3.1 for definition of geographic scope)

- A. Not known from the northeastern US and adjacent Canada 0
- B. Present as a non-native in one northeastern USA state and/or eastern Canadian province. 1
- C. Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces. 2
- D. Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 1 northeastern state or eastern Canadian province. 3
- E. Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces. and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 2 northeastern states or eastern Canadian provinces. 4
- U. Unknown

Score

Documentation:

Identify states and provinces invaded:

KY, IN, IL, IA, MA, MD, MI, MN, MO, NJ, NY, OH, PA, VA, WI, WV; Ont.

Sources of information:

See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.

U.S.D.A. PLANTS database, 2012.

3.6. Current introduced distribution of the species in natural areas in Indiana

- A. Present in no Indiana counties 0
- B. Present in 1-10 Indiana counties 1
- C. Present in 11-20 Indiana counties 2
- D. Present in 21-50 Indiana counties 3
- E. Present in more than 50 Indiana counties or on Federal noxious weed list 4
- U. Unknown

Score

Documentation:

Describe distribution:

See A1.1

Sources of information:

Total Possible
Section Three Total

4. DIFFICULTY OF CONTROL

4.1. Seed banks

- A. Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make viable seeds or persistent propagules. 0
- B. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years 2
- C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years 3
- U. Unknown

Score

INDIANA
NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Documentation:

Identify longevity of seed bank:

Roberts found 0.9% of common teasel seeds to be viable after 5 years storage in the soil (Roberts, 1986). After storage in water for 9 months, only 2% of the seeds germinated successfully and 0% for all seeds stored for longer than 9 months (Comes et al., 1978).

Sources of information:

Comes et al., 1978; Roberts, 1986.

4.2. Vegetative regeneration

- A. No regrowth following removal of aboveground growth 0
- B. Regrowth from ground-level meristems 1
- C. Regrowth from extensive underground system 2
- D. Any plant part is a viable propagule 3
- U. Unknown 3

Score

Documentation:

Describe vegetative response:

Regrowth from basal rosettes.

Sources of information:

Grauver, 2006; Jacquart personal observation.

4.3. Level of effort required

- A. Management is not required: e.g., species does not persist without repeated anthropogenic disturbance. 0
- B. Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft²). 2
- C. Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). 3
- D. Management requires a major investment: e.g. more than 100 person-hours/year of manual effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. Eradication may be impossible (infestation as above). 4
- U. Unknown 4

Score

Documentation:

Identify types of control methods and time-term required:

The following is from Grouver (2006): "The Invasive Plant Association of Wisconsin (IPAW) regards this species as relatively difficult to control (IPAW 2003). Mechanical control is recommended in natural areas. In small stands, rosettes can be dug up, although plants often resprout if the root is not completely removed and damage to the surrounding area can occur if plants are large. Stalks can also be cut once flowering has begun, but before seed set. Because seeds can develop on immature heads, however, the cut stalks need to be removed from the area. Also, cutting of flowering stems may need to be repeated for several years to achieve effective control. Mowing is not an effective control, and in fact often increases the size of patches (Parrish et al. 2005). If mechanical control is not feasible, foliar application of herbicides can be used. Because rosettes of this species are green in early spring and late fall when many native plants are dormant, herbicide control during these times will minimize damage to native species. Also, dicot-selective herbicides (e.g. Triclopyr) are effective, which reduces damage to native monocots. As with mechanical

INDIANA

NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
 Form originally created for use in New York
 Indiana Form version date: November 1, 2010

control, however, herbicide applications over several years are required to manage an established population. Periodic prescribed burning may be helpful in conjunction with mechanical and/or chemical control (Glass, 1990; Weber, 2003; Smith, 2004; WIDNR, 2004; Czarapata 2005). No biocontrol agents are currently in use, but these are being researched (Rector et al. 2006).

"Several years (up to 5-6) of treatment may be necessary to totally eradicate this species from a natural community, regardless of whether mechanical or chemical treatment is chosen (Glass, 1990; Gremaud & Smith, 2002; Smith, 2004; WIDNR, 2004; Czarapata, 2005).

"If rosettes are dug up, damage to the surrounding area can occur if plants are large. If flowering stems are cut, native species of similar height may also be cut in the process. If herbicides are used, non-target damage may occur, though this can be minimized by spraying during the dormant season and/or using a dicot-specific herbicide (Glass, 1990; Weber, 2003; Smith, 2004; WIDNR, 2004; Czarapata, 2005)."

Gucker says: "Available literature (as of 2009) suggest that common teasel may be more susceptible to cutting than cut-leaved teasel."

Sources of information:

Fellows, 2004; Gucker, 2009

Total Possible	10
Section Four Total	7

Total for 4 sections Possible	90
Total for 4 sections	65

References for species assessment:

Brooklyn Botanic Garden. 2009. AILANTHUS database. [Accessed on Oct. 22, 2009].

Comes, R. D.; V. F. Bruns.; & A. D. Kelley. 1978. Longevity of certain weed and crop seeds in fresh water. *Weed Science*. 26(4): 336-344.

Czarapata, E. J. 2005. *Invasive Plants of the Upper Midwest*. The University of Wisconsin Press. Madison, WI. 215 pp.

Fellows, M. & K. Gravuer (rev.) 2006. *Dipsacus fullonum*. U.S. Invasive Species Impact Rank (I-Rank). NatureServe Explorer. <www.natureserve.org>. [Accessed on Aug. 26, 2012].

Donaldson, S. & D. Rafferty. 2002. Identification and management of common teasel (*Dipsacus fullonum*). Fact Sheet-02-40. Reno, NV: University of Nevada, Cooperative Extension. 2 pp.
 <<http://www.unce.unr.edu/publications/files/nr/2002/FS0240.pdf>>. [Accessed on Aug. 26, 2012].

Glass, W. 1990. Vegetation management guideline: Cut-leaved teasel (*Dipsacus laciniatus* L.), Common teasel (*Dipsacus sylvestris* Huds.). Vol. 1, No. 24. Illinois Nature Preserves Commission.
 <<http://dnr.state.il.us/inpc/pdf/VMG%20Teasels%20revised%202007.pdf>> [Accessed on Aug. 27, 2012].

Gremaud, G. & T. Smith. 2002. Teasel Alert! Common and cut-leaved teasels - two species - one BIG problem! Missouri Department of Conservation. <<http://mdc4.mdc.mo.gov/Documents/173.pdf>>. [Accessed on Aug. 27, 2012].

Gucker, C. L. 2009. *Dipsacus fullonum*, *D. laciniatus*. In: Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory.
 <<http://www.fs.fed.us/database/feis/>>. [Accessed on Aug. 26, 2012].

INDIANA

NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE
Form originally created for use in New York
Indiana Form version date: November 1, 2010

Hubbell, S. P. & P. A. Werner. 1979. On measuring the intrinsic rate of increase of populations with heterogeneous life histories. *The American Naturalist*. 113(2): 277-293.

Invasive Plants Association of Wisconsin (IPAW). 2003. IPAW working list of the invasive plants of Wisconsin: a call for comments and information. *Plants Out of Place*. 4. <<http://www.ipaw.org/newsletters/issue4.pdf>>. [Accessed on Aug. 26, 2012].

Knight, R. L.; D. A. Every.; & A. W. Erickson. 1979. Seasonal food habits of four game bird species in Okanogan County, Washington. *The Murrelet*. 60(2): 58-66.

Mittelbach, G. G. & K. L. Gross. 1984. Experimental studies of seed predation in old-fields. *Oecologia*. 65: 7-13.

Parrish, J.; A. Oliver; R. Wiedenmann; S. Post; C. Helm; & M. Timpe. 2005. Effects of mowing on seed dispersal and patch growth of cut leafed teasel (*Dipsacus laciniatus*). Abstract of poster presented at 2005 Ecological Society of America meeting, Montreal, Canada. <abstracts.co.allenpress.com/pweb/esa2005/document/?ID=51710> [Accessed Aug. 27, 2012].

Roberts, H. A. 1986. Seed persistence in soil and seasonal emergence in plant species from different habitats. *Journal of Applied Ecology*. 23(2): 639-656.

Shaw, P. J. & K. Shackleton. 2011. Carnivory in the Teasel *Dipsacus fullonum* — The Effect of Experimental Feeding on Growth and Seed Set. *PLoS ONE* 6(3): e17935. <<http://www.plosone.org/>>. [Accessed on Aug. 26, 2012].

Smith, T. E. 2004. Missouri vegetation management manual. Missouri Department of Conservation, Jefferson City. <http://mdc.mo.gov/sites/default/files/resources/2010/05/5398_3326.pdf>. [Accessed on Aug. 27, 2012].

Snyder, David; Kaufman, Sylvan R. 2004. An overview of nonindigenous plant species in New Jersey. Trenton, NJ: New Jersey Department of Environmental Protection, Division of Parks and Forestry, Office of Natural Lands Management, Natural Heritage Program. 107 pp. <<http://www.nj.gov/dep/parksandforests/natural/heritage/InvasiveReport.pdf>>. [Accessed on Aug. 26, 2012].

United States Department of Agriculture. 2012. Natural Resources Conservation Service. PLANTS database. <<http://plants.usda.gov/java/profile?symbol=DIFU2>> [Accessed on Aug. 26, 2012]

Uva, R. H.; J. C. Neal; & J. M. DiTomaso., eds. 1997; as cited by Gucker, 2009. *Weeds of the Northeast*. New York: Cornell University Press. 397 pp.

Weber, E. 2003. *Invasive plant species of the world: a reference guide to environmental weeds*. CABI Publishing, Cambridge, Massachusetts. 548 pp.

Werner, P. A. 1972; as cited by Werner, 1975. Effect of the invasion of *Dipsacus sylvestris* on plant communities in early old-field succession. Ph.D. thesis, Michigan State University. 140 pp. University Microfilms. Ann Arbor. Mich.

Werner, P. A. 1975. The biology of Canadian weeds. 12. *Dipsacus sylvestris* Huds. *Canadian Journal of Plant Science*. 55: 783-794.

Wisconsin Department of Natural Resources (WIDNR). 2004, September 3 last update. Cut-leaved teasel (*Dipsacus laciniatus*). Invasive species fact sheet. <http://dnr.wi.gov/topic/invasives/fact/commontesael.html> [Accessed. 2012].

INDIANA

NON-NATIVE PLANT INVASIVENESS RANKING FORM

ASSESSMENT FOR INVASIVE PLANTS NOT IN TRADE

Form originally created for use in New York

Indiana Form version date: November 1, 2010

Citation: This IN ranking form may be cited as: Jacquart, E.M. and P.M.Paulone. 2011. Invasiveness ranking system for non-native plants of Indiana. Unpublished. Invasive Plant Advisory Committee (IPAC) to the Indiana Invasive Species Council, Indianapolis, IN.

Acknowledgments: The IN ranking form is an adaptation for Indiana use of the form created for New York by Jordan et al. (2009), cited below. Documentation for species assessed for New York are used for Indiana where they are applicable. The Invasive Plant Advisory Committee was created by the Indiana Invasive Species Council in October 2010, and is made up of the original members of the Indiana Invasive Plant Assessment Working Group (IPSAWG). Original members of IPSAWG included representatives of the The Nature Conservancy; Indiana Native Plant and Wildflower Society; Indiana Nursery and Landscape Association; Indiana Chapter of the American Society of Landscape Architects; Indiana Forage Council; Indiana Wildlife Federation; Indiana State Beekeepers Association; Indiana Beekeeper's Association; Department of Natural Resources; Hoosier National Forest; Indiana Academy of Science; Natural Resources Conservation Service; Indiana Department of Environmental Management; Indiana Department of Transportation; Purdue Cooperative Extension Service; Seed Administrator, Office of the Indiana State Chemist.

References for the Indiana ranking form:

Jordan, M.J., G. Moore, and T.W. Weldy. 2009. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY.

References for the New York ranking form:

Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: http://akweeds.uaa.alaska.edu/akweeds_ranking_page.htm.

Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).

Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. <http://www.natureserve.org/getData/plantData.jsp>

Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. *Invasive Plant Science and Management* 1:36–49

Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M. Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at www.caleppc.org and www.swvma.org. California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.

Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. *Science for Conservation* 209. New Zealand Department of Conservation. 1-23 pp.