

**Assessment of Invasive Species in
Indiana's Natural Areas**

*****OFFICIAL Tall Fescue (*Schenodorus arundinacea*) ASSESSMENT*****

Answers are underlined and in **bold**, comments are inserted in *italics*

Assessed at September 1, 2004 meeting by Ellen Jacquart (TNC), Richard Phillabaum (INDOT), Tom Good (Div. of Reclamation), Keith Johnson (Purdue Univ. Forage), Jennifer Rudgers (IU), Phil O'Connor (Div. of Forestry), Keith Clay (IU), Bob Waltz (Div. of Entomology and Plant Pathology), Larry Nees (State Chemist and Seed Commissioner), Kenneth Smith (Div. of Water), Darrell Brown (NRCS Forage), Lee Casebere (Div. of Nature Preserves), Dan Saffen (Div. of Engineering), Brian MacGowan (Wildlife Society), Gary Langell (Div. of Fish and Wildlife)

Invasive Ranking Summary	Score	
Ecological Impacts	10.5	
Potential For Expansion	31	
Difficulty of Management	23	
Total Score:	64.5	Medium
<i>Rankings: Low < 45, Medium 45 – 80, High > 80</i>		

Contents of the Assessment:

Section I – Invasion Status. Determines whether the species being evaluated is invasive in Indiana.

Section II – Ecological Impacts of Invasion. Evaluates the significance of impacts of the species.

Section III – Potential for Expansion. Evaluates the actual and/or potential expansion of the species.

Section IV – Difficulty of Management. Evaluates how hard it is to control the invasive species.

Section V – Commercial Value. Evaluates how valuable the species is economically in Indiana.

Questions in Sections I – V may direct you to one or more of the following sections for particular invasive species:

Section A. For species which have impacts limited to a few sites, assesses the potential for further spread.

Section B. For species which have medium impacts but high value, assesses whether species could be used in specific circumstances that would prevent escape and invasion.

A worksheet for use with the assessment is found on page 9.

Automatic Exemption from the Assessment

Is this species listed on any federal or on an Indiana state noxious, or prohibited plant lists?

If **YES** then do not proceed with assessment but indicate a conclusion of

Do not use this plant on the front of the response form.

If **NO** then go to Section I.

Section I

Invasion Status

1-a Current Invasion in Indiana

1. Does this species occur in any natural areas in Indiana?

If **NO** then go to Section III-c.

If **YES** then go to 1-a 2.

2. Does it **ONLY** occur in natural areas of Indiana because it has persisted from its previous cultivation

(e.g., in abandoned farmland or homesteads)?

If **YES** then go to Section III-c.

If **NO** then go to Section 1-b (below).


There was a lot of discussion on this. Most agreed that tall fescue could be found in many natural areas, but it may be from past plantings or nearby populations. We had no direct evidence for movement of this species out of plantings and into natural areas, but there are at least a few cases where it seems very unlikely that the fescue population is from a planting. All agreed it is tremendously persistent and does hamper efforts to restore fields to trees or prairie.

1-b Invasion Status in Indiana

Evidence of invasion (forming self-sustaining and expanding populations within a plant community with which it had not previously been associated) must be provided. If not available in a published, quantitative form, this evidence must include written observations from at least three appropriate biologists.

1. Is species invasive **ONLY** when natural disturbance regime and scale have been altered? (e.g. where frequency, extent, or severity of fires have been reduced by human activity).
If **YES** then go to questions 1-b 2.
If **NO** – the species is invasive, go to Section II (below).
2. Has this species ever been known to persist, following colonization, when the natural regime is resumed and the natural flora/communities recover? (e.g., is not an early successional species that only temporarily invades disturbed sites.)
If **YES** (or unknown) - the species is invasive, go to Section II (below).
If **NO** (known not to persist) the species is currently not invasive in Indiana. Go to Section III-c to assess the species' potential for future invasion.

Reported Status of Tall Fescue, *Schenodorus arundinacea*



Section II

Ecological Impacts of Invasion

Impact Index

II-a **Known Impacts at WORST SITE(S) (without, or before, any control effort)**

Add up points for ALL impact statements (i through vi) that are true at the worst affected site(s) then go to question II-b. Evidence of impacts must be provided. If not available in published, quantitative form, this evidence must include written observations from at least *three* appropriate biologists, including specific locations of observations. Scientific names of impacted species (e.g., State-listed or native species with which hybridization occurs) must be included on the response form. If there is no evidence of an impact, then assign 0 points unless the impact is considered very likely (e.g., fixes N₂ in low nutrient soil that can change the flora) or the impact (except vi) has been demonstrated in similar habitats in states. In these cases assign 0.5 points.

i) Causes long-term, broad alterations in ecosystem processes changing the

Points

community as a whole (e.g. invasion of cattails changes hydrology, drying the site and allowing open aquatic systems to become forested). 15

- ii) Has negatively impacted Indiana State-listed or Federal-listed plants or animals (choose one of the following):

Displacement, death or hybridization has been documented AND occurs in at least 20% of known locations of the listed species, OR these effects occur in less than 20% of known locations of the listed species, but at least 4 different listed species are affected. 12

Displacement, death or hybridization occurs in less than 20% of locations of the listed species OR impacts are considered likely because the listed and invasive species closely co-habit (e.g., compete for light). 4

- iii) Displaces or precludes native vegetation (affecting mortality and/or recruitment) by achieving infestations in the state that have at least 50% coverage of this species (as defined in the glossary) in the affected stratum that meet any of the following criteria:

- a) collectively add up to at least 10 acres
- b) are 5 infestations of at least 0.25 acres
- c) are 5 infestations that cover an entire localized community (e.g. sinkhole, seeps, fens, bogs, barrens, cliffs)
- d) are 5 infestations some of which are at least 0.25 acres and others of which cover entire localized communities. 12

A number of studies have shown that tall fescue hinder diversity and growth of some species of plants (Batcher, 2000) but it is not clear that there are enough infestations in natural areas to meet any one of these choices.

- iv) Changes community structure in ways other than vegetation displacement (e.g., alters wildlife abundance, adds a new stratum, or increases stem density within a stratum by more than 5-fold). 4

Batcher (2000): "Songbirds consume tall fescue seeds, and both seeds and foliage are used by small mammals (Wasser 1982). However, some small mammals feeding on fescue infected by the endophyte may become ill as a result of the toxins the endophyte produces (see Impacts and Threats Posed by Tall Fescue section above). Fields dominated by F. arundinacea were found to be poor habitat for bobwhite and quail in Kentucky due to lack of high quality, preferred foods, and improper vegetation structure and composition for nesting and foraging habitat (Barnes et al. 1994). Tall fescue palatability for elk has been reported as poor (Wasser 1982) and elk may show a preference for other grasses. Several studies showed varying preferences by white tailed deer for F. arundinacea (Fire Effects Information System (1996))." Because of its ability to outcompete native vegetation, tall fescue should not be used for wetland mitigation, reforestation, or rehabilitation where managing for wildlife and plant diversity are intended (Burchick 1993)."

Sources of information:

Wasser, 1982; Barnes et al., 1994; FEIS, 1996; Batcher, 2000

- v) Hybridizes with native Indiana plants or commercially-available species. 4

- vi) Covers over 15% of invaded stratum (but if 12 points were assigned for statement iii, do not assign points here) on > 10 acres in the state. 3

Total points (place in worksheet page 9): 7

II-b Range of Habitats in Which Species is Invasive

- Forest: 1)Dry upland, 2)Dry-mesic upland, 3)Mesic upland, 4)Mesic floodplain, 5)Wet-mesic floodplain, 6)Wet floodplain, 7)Bluegrass till plain flatwoods*, 8)Boreal flatwoods*, 9)Central till plain flatwoods, 10)Dry flatwoods*, 11)Sand flatwoods*, 12)Southwestern lowland mesic flatwoods*
- Savanna: 13)Mesic savanna*, 14)Dry sand savanna*, 15)Dry-mesic sand savanna*
- Barrens: 16)Limestone bedrock*, 17)Sandstone bedrock*, 18)Siltstone bedrock*, 19)Chert*, 20)Gravel*, 21)Sand*, 22) Clay*
- Prairie: 23)Dry-mesic prairie*, 24)Mesic prairie*, 25)Wet prairie*, 26)Dry sand prairie*, 27)Dry-mesic sand prairie*, 28)Wet-mesic sand prairie*, **29)Wet sand prairie***
- Wetland: **30)Marl beach*, 31)Acid bog*, 32)Circumneutral bog*, 33)Fen*, 34)Forested fen*, 35)Muck and Sand flats*, 36)Marsh, 37)Sedge meadow*, 38)Panne*, 39)Acid seep*, 40)Calcareous seep*, 41)Circumneutral seep*, 42)Forest swamp, 43)Shrub swamp**
- Lake: **44)Lake, 45)Pond**
- Stream: **46)Low-gradient creek, 47)Medium-gradient creek, 48)High-gradient creek, 49)Low-gradient river, 50)Medium-gradient river, 51)Major river**
- Primary: **52)Aquatic cave*,** 53)Terrestrial cave*, 54)Eroding cliff*, 55)Limestone cliff*, 56)Overhang cliff*, 57)Sandstone cliff*, 58)Lake dune*, 59)Gravel wash*

Is this species known to be invasive in at least four habitat-types (note – rare habitat-types are marked with a * and count as 2 when adding) OR does it occur in at least one habitat-type of each of the terrestrial and palustrine/aquatic lists (palustrine/aquatic habitats are shown in **bold**) 6 non-rare plus 4 rare = 14

If YES then multiply total score from II-a by 1.5
then go to Section II-c (Below)
If **NO** then multiply total score from II-a by 1
then go to Section II-c (Below)

Place point total in worksheet, page 9.

II-c Proportion of Invaded Sites with Significant Impacts

Of the invaded sites, might any of the worst impacts [items i-v in section II-a] only occur under a few, identifiable, environmental conditions (i.e., edaphic or other biological conditions occurring in 1-10% of the sites)? Documentation of evidence must be provided for a **YES** answer.

If **NO** or **NO SCORE** on items i to v in section II-a
then go to Section III
If **YES** then go to Section A

Section III	Potential for Expansion	Potential Index
--------------------	--------------------------------	------------------------

This section evaluates a species’ actual and/or potential for expansion in Indiana.

III-a Potential for Becoming Invasive in Indiana

1. Is information available on the occurrence of new populations of this species in Indiana over the last 5 years?

If **YES** then go to section III-b

If **NO** go to Section III-c to estimate potential for expansion based on the biology of the species.

III-b. Known Rate of Invasion.

1. Was this species reported in more than two new discrete sites (e.g., lakes, parks, fragments of habitats at least 5 miles apart) in any 12 month period within the last 5 years?
 - If **NO** then P = Low; then go to Section IV
 - If **YES** then P = High; then go to Section IV

III-c. Estimated Rate of Invasion. This section is used to predict the risk of invasion for species that are 1) not currently invasive in the state, and 2) invasive in the state but for which no data on current rate of spread exists. These questions are based on Hiebert et al. 1995.

1. Does this species hybridize with any State-listed plants or commercially-important species? (E.g., exhibit pollen / genetic invasion.)
 - If **YES** then go to Section B
 - If **NO** then go to question III-c 2.
 - It may rarely hybridize with *Lolium perenne*.*

2. Add up all points from statements that are true for this species. Points

- i. Ability to complete reproductive cycle in area of concern
 - a. not observed to complete reproductive cycle 0
 - b. observed to complete reproductive cycle 5

- ii. Mode of reproduction
 - a. reproduces almost entirely by vegetative means 1
 - b. reproduces only by seeds 3
 - c. reproduces vegetatively and by seed 5

Reproduces by seed and spreads vegetatively (Batcher, 2000).

- iii. Vegetative reproduction
 - a. no vegetative reproduction 0
 - b. vegetative reproduction rate maintains population 1
 - c. vegetative reproduction rate results in moderate increase in population size 3
 - d. vegetative reproduction rate results in rapid increase in population size 5

- iv. Frequency of sexual reproduction for mature plant
 - a. almost never reproduces sexually in area 0
 - b. once every five or more years 1
 - c. every other year 3
 - d. one or more times a year 5

- v. Number of seeds per plant
 - a. few (0-10) 1
 - b. moderate (11-1,000) 3
 - c. many-seeded (> 1,000) 5

- vi. Dispersal ability
 - a. little potential for long-distance dispersal 0
 - b. great potential for long-distance dispersal 5

Seeds remain viable after passing through the digestive track of animals (Burchick, 1993 as cited in Batcher, 2000).

vii. Germination requirements

- a. requires open soil and disturbance to germinate 0
- b. can germinate in vegetated areas but in a narrow range or in special conditions 3
- c. can germinate in existing vegetation in a wide range of conditions 5

Batcher notes: “[Wolf et al., 1979] found that the minimum temperature for germination is 4.5°C; optimum ranges of both 12 to 18°C and 20-25°C have been found” (as cited in Batcher, 2000). Tall fescue seeds sown the year of stored in uncontrolled environments (Idaho and Nebraska) for 19 years were found to have a germination rate of 4.5%; seeds sown from the same year germinated at a rate of 93% (Hull, 1973).

viii. Competitive ability

- a. poor competitor for limiting factors 0
- b. moderately competitive for limiting factors 3
- c. highly competitive for limiting factors 5

*Used specially for acid mine spoil reclamation and is highly tolerant of low pH (4-6) and manganese concentrations (4-64 ppm) (Fleming et al. 1974). Burchick recommends *F. arundinacea* to not be planted in wetlands (Burchick, 1993). It was noted by Wheeler & Hill that: “[*F. arundinacea*] colonizes bare soil, and is a strong competitor in many species mixtures” (as cited in Batcher, 2000).*

Total points for questions i – viii (place in worksheet page 9): 31

Section IV	Difficulty of Management	Management Index
------------	--------------------------	------------------

IV Factors That Increase the Difficulty of Management

Add up all points from statements that are true for this species then go to Section V. Assign 0.5 point for each statement for which a true/false response is not known.

	<u>Points</u>
i) Control techniques that would eliminate the worst-case effects (as listed in Section II) have been investigated but none has been found.	15
ii) This species is difficult to control without significant damage to native species because: it is widely dispersed throughout the sites (i.e., does not occur within discrete clumps nor monocultures); it is attached to native species (e.g., vine, epiphytes or parasite); or there is a native plant which is easily mistaken for this invader in: (choose one)	
≥ 50% of discrete sites in which this species grows;	10
25% to 50% of discrete sites in which this species grows.	7
iii) Total contractual costs of known control method per acre in first year, including access, personnel, equipment, and materials (any needed re-vegetation is not included) > \$2,000/acre (estimated control costs are for acres with a 50% infestation)	5
iv) Further site restoration is <i>usually</i> necessary following plant control to reverse ecosystem impacts and to restore the original habitat-type or to prevent immediate re-colonization of the invader.	<u>5</u>
v) The total area over which management would have to be conducted is: (choose one)	
≥ 100 acres;	<u>5</u>

< 100 but > 50 acres.	2
≤ 50 but > 10 acres.	1
≤10 acres	1/2
vi) Following the first year of control of this species, it would be expected that individual sites would require re-survey or re-treatment, due to recruitment from persistent seeds, spores, or vegetative structures, or by dispersal from outside the site: (choose one)	
at least once a year for the next 5 years;	<u>10</u>
one to 4 times over the next 5 years;	6
regrowth not known	2
vii) Occurs in more than 20 discrete sites (e.g., water-basins, parks, fragments of habitats at least 5 miles apart).	<u>3</u>
viii) The number of viable, independent propagules per mature plant (e.g., seeds, spores, fragments, tubers, etc. detached from parent) is > 200 per year AND one or more of the following:	
A. the propagules can survive for more than 1 year;	
B. the propagules have structures (fleshy coverings, barbs, plumes, or bladders) that indicate they may spread widely by birds, mammals, wind or water;	
C. the infestations at 3 or more sites exhibit signs of long distance dispersal. Some possible indicators of long distance dispersal include: the infestation has outlier individuals distant [>50 yards] from the core population; the infestation apparently lacks sources of propagules within ¼ mile.	3
ix) Age at first reproduction is within first 10% of likely life-span and/or less than 3 months.	2
Total points (place in worksheet page 9):	23

Section V	Commercial Value	Value Index
------------------	-------------------------	--------------------

V-a Commercial Value

Does this species have any commercial value?

If response is **NO** then V = 0 and Go to Conversion of Index Scores to Index Categories

If response is **YES** then go to Section V-b

V-b Factors that Indicate a Significant Commercial Value

Larry Nees brought information on the sale of tall fescue in the state. There are roughly 26 distributors in the state and many retail outlets. Of the 50-60 retail outlets they have visited this year, approximately 780,000 lbs. of tall fescue seed were sold. Many different varieties were being sold, mostly of the turf type. Total sales of fescue seed in the state are estimated to be at least 1 million pounds/year. INDOT probably accounts for about half of those sales. There has been an increase in the amount of turf-type fescue sold in recent years.

Add up all points from statements that are true for this species. Assign 0.5 point for each statement for which a true/false response is not known.

- | | |
|--|-----------|
| i) This species is sold in national or regional retail stores (e.g., Wal-Mart, Home Depot, Publix). | <u>10</u> |
| ii) State-wide there are more than 20 commercial growers of this species.
<i>For sod.</i> | <u>7</u> |
| iii) More than five growers in Indiana rely on this species as more than 10% of their nursery's production.
<i>For sod.</i> | <u>3</u> |
| iv) This species has provided a crop, turf, or feed source (e.g., forage, nectar) that has been, or resulted in, a significant source of income for at least five farmers for over 20 years. | <u>3</u> |
| v) This species is utilized statewide | <u>3</u> |
| vi) There are more than 100 retail seed outlets statewide | <u>3</u> |

Total points (place in worksheet page 9): **29**

Section A (from Section II-c)

A1 Can the habitats in which the worst-case ecological impacts occur (items i to v in Section II-a) be clearly defined as different from invaded sites where there are no such impacts (e.g., defined by edaphic or biological factors)? (If ecological impacts include negative effects on a State-listed species, then the specific habitats in which that State-listed species occurs must be clearly distinguishable from habitats in which it does not occur.)

If **NO** then return to Section III

If **YES** then go to question A2 and prepare such a site definition

A2 Can an estimate be made of the maximum distance that propagules (or pollen if hybridization is a concern) might reasonably be expected to disperse?

If **NO** then return to Section III

If **YES** then prepare instructions for Specified and Limited Use based on maximum dispersal distance (e.g., may be acceptable for use in specific areas but not near habitats where impacts are high.) Reassess if the incidence of worst-case impacts increases above 10% or within 10 years, whichever is earlier. THEN resume the assessment at Section III to provide scores for the other indices.

Section B (from Section III-c or if Value = High and Impact = Medium)

B1 Are there specific circumstances in which this species could be used that would not be expected to result in escape and invasion? (E.g., foliage plants that are only used indoors and which can be reasonably prevented, by conspicuous labeling, from use or disposal in the landscape.)

If **NO**, then retain the previously derived Conclusion.

If **YES**, then Acceptable for Specified and Limited Use where regulations and educational programs for penalties and enforcement of misuse exist. Reassess this species every 2 years.

Worksheet for Assessment

Section I:

Follow directions to different sections.

Section II:Impacts Point Total: 7 X (1.5) = 10.5 **Impacts****Section III:**Potential = High Medium or Low 31 **Potential for Expansion****Section IV:**Difficulty of Management Point Total: 23 **Difficulty of Management****Section V:**Commercial Value Point Total: 29 **Value**

Invasive Ranking Summary:

Invasive Ranking Summary	Score
Ecological Impacts	10.5
Potential For Expansion	31
Difficulty of Management	23
Total Score:	64.5
	Medium
<i>Rankings: Low < 45, Medium 45 – 80, High > 80</i>	

Recommendation

Plant with caution – plant only for high flow grassy waterways and to stabilize slopes steeper than 4:1; avoid using for forage but use low-endophyte forms whenever used.

Literature Cited

- Ball, D.M., J.F. Pedersen, and G.D. Lacefield. 1993. The tall-fescue endophyte. *American Scientist*. 81: 370-379.
- Batcher, M. 2000. Element Stewardship Abstract for *Festuca arundinacea* (Schreb.) Synonym: *Festuca elatior* L. The Nature Conservancy. < <http://www.imapinvasives.org/GIST/ESA/esapages/documnts/festaru.pdf>>. [Accessed on May 23, 2012.]
- Burchick, M. 1993. The problems with tall fescue in ecological restoration. *Wetland Journal*. 5(2):16.
- Fleming, A.L., J.W. Schwartz, and C.D. Foy. 1974. Chemical factors controlling the adaptation of weeping lovegrass and tall fescue to acid mine spoils. *Agronomy Journal*. 66(6): 715-719.
- Henson, J. F. 2001. Tall Fescue *Lolium arundinaceum* (Schreb.) S. J. Darbyshire. NRCS Plant Guide. National Plant Data Center, Baton Rouge, Louisiana.
- Hull, A.C., Jr. 1973. Germination of range plant seeds after long periods of uncontrolled storage. *Journal of Range Management*. 26(3): 198-200.
- Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York: *Festuca arundinacea*. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Revised by Dong Lee, Angela Sturdevant, Ellen Jacquart 9/21/2012

Brooklyn, NY; The Nature Conservancy, Albany, NY

Wheeler, W.A. and D.D. Hill. 1957. Grassland seeds. Van Nostrand Co. Princeton, New Jersey.

Wolf, D.D., R.H. Brown, and R.E. Blaser. 1979. Physiology of Growth and Development. In

Batcher, M. 2000. Element Stewardship Abstract for *Festuca arundinacea* (Schreb.) Synonym: *Festuca elatior* L. The Nature Conservancy

Glossary

Anthropogenic disturbance. Human-induced disturbance (e.g., mowing) or human-induced changes in natural disturbance regime (e.g., changing the frequency, extent, or severity of fires).

Coverage. Visual or quantitative estimate of the relative amount of area in a stratum where the canopy of the non-native species intercepts the light that would otherwise be available for other species in or below that stratum. Estimated cover may be dispersed or continuous in a site. Cover is usually measured when foliage is fully expanded. In the case of species that form a dense, continuous mat of rhizomes or stolons, the percent of the soil surface or upper level occupied by that root mat can be estimated as soil, rather than canopy, cover.

Disturbance. Mechanisms that limit biomass by causing its partial or total destruction.

Discrete sites. Disjunct habitat-types or fragments of habitats at least 1 mile apart that support invasive plant populations that likely arose by separate long-distance dispersal mechanisms.

Federal- or Indiana -listed. Species that are listed by Federal laws or Indiana statutes or rules as threatened or endangered within the State of Indiana. This list with notes is available at <http://www.state.in.us/dnr/naturepr/endanger/plant.htm>

Invasive. A species that forms self-sustaining and expanding populations within a natural plant community with which it had not previously been associated (Vitousek *et al.* 1995).

Long-term alterations in ecosystem processes. Examples of ecosystem processes that could be altered: erosion and sedimentation rates; land elevation; water channels; water-holding capacity; water-table depth; surface flow patterns; rates of nutrient mineralization or immobilization; soil or water chemistry; and type, frequency, intensity, or duration of disturbance. For further explanation see Gordon (1998).

Native. Species within its natural range or natural zone of dispersal (i.e., within the range it could have, or would have, occupied without direct or indirect introduction and/or care by humans. Excludes species descended from domesticated ancestors) (Vitousek *et al.* 1995).

Natural areas. Natural areas: Areas with native plant communities supporting native plant and animal species, with long undisturbed soil systems, and hydrological regimes relatively intact or under restoration. Edges of historically or currently disturbed areas (roadsides, trails, adjacent to historically disturbed locations, etc.) should not be included in the assessment of invasion into natural areas. That invasion may have been facilitated by the edges, but has to have extended into the native communities for inclusion in this category.

Pollen or genetic invasion. When a native species is displaced by a non-native species through hybridization.

Stratum. A distinct layer in the architecture of vegetation (e.g., tree canopy or understory shrubs).