

**Assessment of Invasive Species in
Indiana's Natural Areas**

*****OFFICIAL Common Periwinkle and Big Leaf Periwinkle (*Vinca minor* and *major*)
ASSESSMENT*****

Answers are underlined and in **bold**, comments are inserted in *italics*

Assessed November 5, 2004 by Ken Collins (NRCS, group leader), Ellen Jacquart (TNC), Cliff Chapman (DNR – DNP), Phil O'Connor (DNR – DoF), Mike Cline (INLA), Dave Gorden (ASLA), Lori Johnson (Indy Zoo).

<i>Vinca minor</i>	Score	
Ecological Impacts	42	
Potential for Expansion	10	
Difficulty of Management	19	
Total Score:	71	Medium
<i>Rankings: Low < 45, Medium 45 – 80, High > 81</i>		

<i>Vinca major</i>	Score	
Ecological Impacts	0	
Potential for Expansion	0	
Difficulty of Management	0	
Total Score:	0	Caution*
<i>Rankings: Low < 45, Medium 45 – 80, High > 81</i>		

***Note** – *Vinca major*, big leaf periwinkle, was also discussed as part of this assessment. Currently, it is known to have spread from plantings in 9 counties in Indiana (per Overlease and Yatskievych); however, the spread is minor (no more than 25 sq. ft. in most cases) and limited to disturbed areas. We felt this did not meet the definition of ‘invasive’ and so did not complete the assessment for this species. We do feel this species should be watched, as it has become invasive in some parts of the country. It will be given a ‘caution’ ranking.

Contents of the Assessment:

- Section I** – Invasion Status. Determines whether the species being evaluated is invasive in Indiana.
- Section II** – Ecological Impacts of Invasion. Evaluates the significance of impacts of the species.
- Section III** – Potential for Expansion. Evaluates the actual and/or potential expansion of the species.
- Section IV** – Difficulty of Management. Evaluates how hard it is to control the invasive species.
- Section V** – Commercial Value. Evaluates how valuable the species is economically in Indiana.

Questions in Sections I – V may direct you to one or more of the following sections for particular invasive species:

- Section A.** For species which have impacts limited to a few sites, assesses the potential for further spread.
- Section B.** For species which have medium impacts but high value, assesses whether species could be used in specific circumstances that would prevent escape and invasion.

A worksheet for use with the assessment is found on page 9.

Automatic Exemption From the Assessment

Is this species listed on any federal or on an Indiana state noxious, or prohibited plant lists?

If **YES** then do not proceed with assessment but indicate a conclusion of
Do not use this plant on the front of the response form.

If **NO** then go to Section I.

Section I

Invasion Status

1-a Current Invasion in Indiana

1. Does this species occur in any natural areas in Indiana?

If **NO** then go to Section III-c.

If **YES** then go to 1-a 2.

2. Does it **ONLY** occur in natural areas of Indiana because it has persisted from its previous cultivation (e.g., in abandoned farmland or homesteads)? *Vinca persists where it is planted, but also spreads vegetatively from the cultivated site into undisturbed natural areas.*

If **YES** then go to Section III-c.

If **NO** then go to Section 1-b (below).

1-b Invasion Status in Indiana

Evidence of invasion (forming self-sustaining and expanding populations within a plant community with which it had not previously been associated) must be provided. If not available in a published, quantitative form, this evidence must include written observations from at least three appropriate biologists.

1. Is species invasive **ONLY** when natural disturbance regime and scale have been altered? (e.g. where frequency, extent, or severity of fires have been reduced by human activity).

If **YES** then go to questions 1-b 2.


If **NO** – the species is invasive, go to Section II (below).

2. Has this species ever been known to persist, following colonization, when the natural regime is resumed and the natural flora/communities recover? (e.g., is not an early successional species that only temporarily invades disturbed sites.)


If **YES** (or unknown) - the species is invasive, go to Section II (below).

If **NO** (known not to persist) the species is currently not invasive in Indiana. Go to Section III-c to assess the species' potential for future invasion.

Reported Status of Bigleaf Periwinkle, *Vinca Major*



Reported Status of Common Periwinkle, *Vinca minor*



Section II

Ecological Impacts of Invasion

Impact Index

II-a Known Impacts at WORST SITE(S) (without, or before, any control effort)

Add up points for ALL impact statements (i through vi) that are true at the worst affected site(s) then go to question II-b. Evidence of impacts must be provided. If not available in published, quantitative form, this evidence must include written observations from at least *three* appropriate biologists, including specific locations of observations. Scientific names of impacted species (e.g., State-listed or native species with which hybridization occurs) must be included on the response form. If there is no evidence of an impact, then assign 0 points unless the impact is considered very likely (e.g., fixes N₂ in low nutrient soil that can change the flora) or the impact (except vi) has been demonstrated in similar habitats in states. In these cases assign 0.5 points.

- i) Causes long-term, broad alterations in ecosystem processes changing the community as a whole (e.g. invasion of cattails changes hydrology, drying the site and allowing open aquatic systems to become forested).
Gravuer has cited several references (Drewitz, 2000; Weber 2003; Holloran et al. 2004; Makings, 2005) where dense mats of periwinkle may change erosion processes by displacement of native shrubs and trees in riparian areas (as cited in Gravuer, 2007).

Points

15

- ii) Has negatively impacted Indiana State-listed or Federal-listed plants or

animals (choose one of the following):

Displacement, death or hybridization has been documented AND occurs in at least 20% of known locations of the listed species, OR these effects occur in less than 20% of known locations of the listed species, but at least 4 different listed species are affected. *There is one site in the state for Dentaria multifida, and it is being threatened by Vinca. Thus, there is displacement in 100% of the known locations of this listed species.*

12

Displacement, death or hybridization occurs in less than 20% of locations of the listed species OR impacts are considered likely because the listed and invasive species closely co-habit (e.g., compete for light).

4

iii) Displaces or precludes native vegetation (affecting mortality and/or recruitment) by achieving infestations in the state that have at least 50% coverage of this species (as defined in the glossary) in the affected stratum that meet any of the following criteria:

- a) collectively add up to at least 10 acres
- b) are 5 infestations of at least 0.25 acres
- c) are 5 infestations that cover an entire localized community (e.g. sinkhole, seeps, fens, bogs, barrens, cliffs)
- d) are 5 infestations some of which are at least 0.25 acres and others of which cover entire localized communities.

12

iv) Changes community structure in ways other than vegetation displacement (e.g., alters wildlife abundance, adds a new stratum, or increases stem density within a stratum by more than 5-fold).

4

As noted earlier, periwinkle form dense mats (Drewitz, 2000; Swearingen et al. 2002; Weber 2003; Holloran et al. 2004; Makings, 2005; Ma & Moore, 2009). Allelopathic inhibition may contribute to suppression of seedling growth in addition to shading from dense mats (Darcy & Burkhart, 2002).

v) Hybridizes with native Indiana plants or commercially-available species.

4

vi) Covers over 15% of invaded stratum (but if 12 points were assigned for statement iii, do not assign points here) on > 10 acres in the state.

3

Total points (place in worksheet page 9): 28

II-b Range of Habitats in Which Species is Invasive

Forest: 1)Dry upland, 2)Dry-mesic upland, 3)Mesic upland, 4)Mesic floodplain, 5)Wet-mesic floodplain, 6)Wet floodplain, 7)Bluegrass till plain flatwoods*, 8)Boreal flatwoods*, 9)Central till plain flatwoods, 10)Dry flatwoods*, 11)Sand flatwoods*, 12)Southwestern lowland mesic flatwoods*

Savanna: 13)Mesic savanna*, 14)Dry sand savanna*, 15)Dry-mesic sand savanna*

Barrens: 16)Limestone bedrock*, 17)Sandstone bedrock*, 18)Siltstone bedrock*, 19)Chert*, 20)Gravel*, 21)Sand*, 22) Clay*

Prairie: 23)Dry-mesic prairie*, 24)Mesic prairie*, **25)Wet prairie***, 26)Dry sand prairie*, 27)Dry-mesic sand prairie*, 28)Wet-mesic sand prairie*, **29)Wet sand prairie***

Wetland: **30)Marl beach*, 31)Acid bog*, 32)Circumneutral bog*, 33)Fen*, 34)Forested fen*, 35)Muck and Sand flats*, 36)Marsh, 37)Sedge meadow*, 38)Panne*, 39)Acid seep***,

40)Calcareous seep*, 41)Circumneutral seep*, 42)Forest swamp, 43)Shrub swamp

Lake: 44)Lake, 45)Pond

Stream: 46)Low-gradient creek, 47)Medium-gradient creek, 48)High-gradient creek, 49)Low-gradient river, 50)Medium-gradient river, 51)Major river

Primary: 52)Aquatic cave*, 53)Terrestrial cave*, 54)Eroding cliff*, 55)Limestone cliff*, 56)Overhang cliff*, 57)Sandstone cliff*, 58)Lake dune*, 59)Gravel wash*

Is this species known to be invasive in at least four habitat-types (note – rare habitat-types are marked with a * and count as 2 when adding) OR does it occur in at least one habitat-type of each of the terrestrial and palustrine/aquatic lists (palustrine/aquatic habitats are shown in **bold**)

If YES then multiply total score from II-a by 1.5
then go to Section II-c (Below)

If NO then multiply total score from II-a by 1
then go to Section II-c (Below)

Place point total in worksheet, page 9.

II-c Proportion of Invaded Sites with Significant Impacts

Of the invaded sites, might any of the worst impacts [items i-v in section II-a] only occur under a few, identifiable, environmental conditions (i.e., edaphic or other biological conditions occurring in 1-10% of the sites)? Documentation of evidence must be provided for a **YES** answer.

**If NO or NO SCORE on items i to v in section II-a
then go to Section III**

If YES then go to Section A

Section III

Potential for Expansion

Potential Index

This section evaluates a species' actual and/or potential for expansion in Indiana.

III-a Potential for Becoming Invasive in Indiana

1. Is information available on the occurrence of new populations of this species in Indiana over the last 5 years?

If **YES** then go to section III-b

If **NO** go to Section III-c to estimate potential for expansion based on the biology of the species.

III-b. Known Rate of Invasion.

1. Was this species reported in more than two new discrete sites (e.g., lakes, parks, fragments of habitats at least 5 miles apart) in any 12 month period within the last 5 years?

If **NO** then P = Low; then go to Section IV

If **YES** then P = High; then go to Section IV

III-c. Estimated Rate of Invasion. This section is used to predict the risk of invasion for species that are 1) not currently invasive in the state, and 2) invasive in the state but for which no data on current rate of spread exists. These questions are based on Hiebert et al. 1995.

1. Does this species hybridize with any State-listed plants or commercially-important species? (E.g., exhibit pollen / genetic invasion.)

If **YES** then go to Section B

If NO then go to question III-c 2.

	<u>Points</u>
2. Add up all points from statements that are true for this species.	
i. Ability to complete reproductive cycle in area of concern	
a. not observed to complete reproductive cycle	<u>0</u>
b. observed to complete reproductive cycle	5
<i>To our knowledge, V. minor is not producing viable seed in Indiana.</i>	
ii. Mode of reproduction	
a. reproduces almost entirely by vegetative means	<u>1</u>
b. reproduces only by seeds	3
c. reproduces vegetatively and by seed	5
<i>There are sparse reports of periwinkle reproducing by seeds (Miller, 2003; Chess, 2005) but the commonly held view is that periwinkle mainly propagate vegetatively (Swearingen, 2002; Stone 2005, Gravuer, 2007). Randall (cited in Stone, 2005) notes that periwinkle does not propagate by seed outside its native habitat.</i>	
iii. Vegetative reproduction	
a. no vegetative reproduction	0
b. vegetative reproduction rate maintains population	1
c. vegetative reproduction rate results in moderate increase in population size	<u>3</u>
d. vegetative reproduction rate results in rapid increase in population size	5
iv. Frequency of sexual reproduction for mature plant	
a. almost never reproduces sexually in area	<u>0</u>
b. once every five or more years	1
c. every other year	3
d. one or more times a year	5
v. Number of seeds per plant	
a. few (0-10)	<u>1</u>
b. moderate (11-1,000)	3
c. many-seeded (> 1,000)	5
<i>Three to five seeds per cylindrical fruit (Miller, 2003).</i>	
vi. Dispersal ability	
a. little potential for long-distance dispersal	<u>0</u>
b. great potential for long-distance dispersal	5
<i>Possible long-distance dispersal is by water or garden refuse (Gravuer, 2007). In its native habitat, periwinkle is dispersed by ants (Honnay et al., 1999; Jacquemyn et al., 2001).</i>	
vii. Germination requirements	
a. requires open soil and disturbance to germinate	<u>0</u>
b. can germinate in vegetated areas but in a narrow range or in special conditions	3
c. can germinate in existing vegetation in a wide range of conditions	5
viii. Competitive ability	
a. poor competitor for limiting factors	0
b. moderately competitive for limiting factors	3
c. highly competitive for limiting factors	<u>5</u>
Total points for questions i – viii (place in worksheet page 9): <u>10</u>	

IV Factors That Increase the Difficulty of Management

Add up all points from statements that are true for this species then go to Section V. Assign 0.5 point for each statement for which a true/false response is not known. *Cliff Chapman noted that while control of this species is difficult, it is not impossible. He has found foliar spray (8% glyphosate plus Nufilm IR) in March provides the best control while avoiding non-target impacts. Spraying in November may control it as well, but leaf litter cover is substantial and prevents one from making contact with all the plants.*

	<u>Points</u>
i) Control techniques that would eliminate the worst-case effects (as listed in Section II) have been investigated but none has been found.	15
ii) This species is difficult to control without significant damage to native species because: it is widely dispersed throughout the sites (i.e., does not occur within discrete clumps nor monocultures); it is attached to native species (e.g., vine, epiphytes or parasite); or there is a native plant which is easily mistaken for this invader in: (choose one)	
≥ 50% of discrete sites in which this species grows;	10
25% to 50% of discrete sites in which this species grows.	7
iii) Total contractual costs of known control method per acre in first year, including access, personnel, equipment, and materials (any needed re-vegetation is not included) > \$2,000/acre (estimated control costs are for acres with a 50% infestation)	5
iv) Further site restoration is <i>usually</i> necessary following plant control to reverse ecosystem impacts and to restore the original habitat-type or to prevent immediate re-colonization of the invader. <i>Cliff and Ellen provided examples where this was the case – recolonization of Vinca sites seems to be quite slow (at least a few years have gone by with no species moving into treated areas).</i>	<u>5</u>
v) The total area over which management would have to be conducted is: (choose one)	
≥ 100 acres;	<u>5</u>
< 100 but > 50 acres.	2
≤ 50 but > 10 acres.	1
≤10 acres	1/2
vi) Following the first year of control of this species, it would be expected that individual sites would require re-survey or re-treatment, due to recruitment from persistent seeds, spores, or vegetative structures, or by dispersal from outside the site: (choose one)	
at least once a year for the next 5 years;	10
one to 4 times over the next 5 years;	<u>6</u>
regrowth not known	2
vii) Occurs in more than 20 discrete sites (e.g., water-basins, parks, fragments of habitats at least 5 miles apart).	<u>3</u>
viii) The number of viable, independent propagules per mature plant (e.g., seeds, spores, fragments, tubers, etc. detached from parent) is > 200 per year AND one or more of the following:	
A. the propagules can survive for more than 1 year;	

- B. the propagules have structures (fleshy coverings, barbs, plumes, or bladders) that indicate they may spread widely by birds, mammals, wind or water;
- C. the infestations at 3 or more sites exhibit signs of long distance dispersal. Some possible indicators of long distance dispersal include: the infestation has outlier individuals distant [>50 yards] from the core population; the infestation apparently lacks sources of propagules within $\frac{1}{4}$ mile. 3

ix) Age at first reproduction is within first 10% of likely life-span and/or less than 3 months. 2

Total points (place in worksheet page 9): **19**

Section V	Commercial Value	Value Index
------------------	-------------------------	--------------------

V-a **Commercial Value**

Does this species have any commercial value?

If response is **NO** then V = 0 and Go to Conversion of Index Scores to Index Categories

If response is **YES** then go to Section V-b

Mike Cline reported that he talked to 7-8 nurseries and found that growers of Vinca in Indiana sell between \$110,000 to \$170,000. Lots more is sold by wholesalers/retailers who purchase the Vinca to sell. The most common cultivar is cv. Bowles.

V-b **Factors that Indicate a Significant Commercial Value**

Add up all points from statements that are true for this species. Assign 0.5 point for each statement for which a true/false response is not known.

- | | <u>Points</u> |
|--|------------------|
| i) This species is sold in national or regional retail stores (e.g., WalMart, Home Depot, Publix). | <u>10</u> |
| ii) State-wide there are more than 20 commercial growers of this species. | 7 |
| iii) More than five growers in Indiana rely on this species as more than 10% of their nursery's production. | 3 |
| iv) This species has provided a crop, turf, or feed source (e.g., forage, nectar) that has been, or resulted in, a significant source of income for at least five farmers for over 20 years. | 3 |
| v) This species is utilized statewide | <u>3</u> |
| vi) There are more than 100 retail seed outlets statewide | 3 |
| Total points (place in worksheet page 9): <u>13</u> | |

Section A (from Section II-c)

A1 Can the habitats in which the worst-case ecological impacts occur (items i to v in Section II-a) be
Edited by Alison Clements, Margaret David, Dong Lee, and Jacob Krebs, July 5, 2012

clearly defined as different from invaded sites where there are no such impacts (e.g., defined by edaphic or biological factors)? (If ecological impacts include negative effects on a State-listed species, then the specific habitats in which that State-listed species occurs must be clearly distinguishable from habitats in which it does not occur.)

If **NO** then return to Section III

If **YES** then Go to question A2 and prepare such a site definition

A2 Can an estimate be made of the maximum distance that propagules (or pollen if hybridization is a concern) might reasonably be expected to disperse?

If **NO** then return to Section III

If **YES** then prepare instructions for Specified and Limited Use based on maximum dispersal distance (e.g., may be acceptable for use in specific areas but not near habitats where impacts are high.) Reassess if the incidence of worst-case impacts increases above 10% or within 10 years, whichever is earlier. THEN resume the assessment at Section III to provide scores for the other indices.

Section B (from Section III-c or if Value = High and Impact = Medium)

B1 Are there specific circumstances in which this species could be used that would not be expected to result in escape and invasion? (E.g., foliage plants that are only used indoors and which can be reasonably prevented, by conspicuous labeling, from use or disposal in the landscape.)

If **NO**, then retain the previously derived Conclusion.

If **YES**, then Acceptable for Specified and Limited Use where regulations and educational programs for penalties and enforcement of misuse exist. Reassess this species every 2 years.

Worksheet for Assessment

Section I:

Follow directions to different sections.

Section II:

Impacts Point Total: 28 X (1 or 1.5) = 42 **Impacts**

Section III:

Potential = High Medium or Low 10 **Potential for Expansion**

Section IV:

Difficulty of Management Point Total: 19 **Difficulty of Management**

Section V:

Commercial Value Point Total: 13 **Value**

Invasive Ranking Summary:

<i>Vinca minor</i>	Score	
Ecological Impacts	42	
Potential for Expansion	10	
Difficulty of Management	19	
Total Score:	71	Medium
<i>Rankings: Low < 45, Medium 45 – 80, High > 81</i>		

<i>Vinca major</i>	Score	
Ecological Impacts	0	
Potential for Expansion	0	
Difficulty of Management	0	
Total Score:	0	Caution*
<i>Rankings: Low < 45, Medium 45 – 80, High > 81</i>		

***Note** – *Vinca major*, big leaf periwinkle, was also discussed as part of this assessment. Currently, it is known to have spread from plantings in 9 counties in Indiana (per Overlease and Yatskievych); however, the spread is minor (no more than 25 sq. ft. in most cases) and limited to disturbed areas. We felt this did not meet the definition of ‘invasive’ and so did not complete the assessment for this species. We do feel this species should be watched, as it has become invasive in some parts of the country. It will be given a ‘caution’ ranking.

Literature Cited

Bean, C. and M. Russo. rev. 1988. Element Stewardship Abstract for *Vinca major*. The Nature Conservancy. <<http://www.imapinvasives.org/GIST/ESA/esapages/documnts/vincmaj.pdf>>. [Accessed on Jun 5, 2012.]

Chess, K. 2005. *Vinca* seen propagating by seed again (California, USA), [Online]. In: Global Invasive Species Team (GIST) listserv digest #139: Posting #10--September 2005. The Nature Conservancy (Producer). <<http://www.invasive.org/gist/listarch/arch139.html#01>> [Jun 5, 2012]

Darcy, A.J. and Burkart, M.C. 2002. Allelopathic potential of *Vinca minor*, an invasive exotic plant in west Michigan forests. *BIOS* 73:127–132

Drewitz, J. 2000. *Vinca major* L. pg. 326-329 in Bossard, C.C., J.M. Randall, and M. Hoshovsky. (eds.) *Invasive Plants of California's Wildlands*. University of California Press, Berkeley, CA.

Gravuer, K. 2007. *Vinca minor*. U.S. Invasive Species Impact Rank (I-Rank). NatureServe Explorer. <www.natureserve.org>. [Accessed on Jun 5, 2012.]

Holloran, P., A. Mackenzie, S. Farrell, and D. Johnson. 2004. *The weed workers' handbook: A guide to techniques for removing Bay Area invasive plants*. The Watershed Project, California Invasive Plant Council. Richmond, CA.

Honnay, O., M. Hermy., and P. Coppin. 1999. Impact of habitat quality on forest plant species colonization. *Forest Ecology and Management*. 115(2-3): 157-170.

Jacquemyn, H., J. Butaye., and M. Hermy. 2001. Forest plant species richness in small, fragmented mixed deciduous forest patches: the role of area, time and dispersal limitation. *Journal of Biogeography*. 28(6): 801-812.

Ma, J. and G. Moore. 2009. Invasiveness ranking system for non-native plants of New York: *Vinca Minor*. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY.

Makings, E. 2005. Plant Assessment Form, for use with "Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands" by the California Exotic Pest Plant Council and the Southwest Vegetation Management Association: *Vinca major* L. <http://sbsc.wr.usgs.gov/research/projects/swepic/SWVMA/PLANTPDF/Vinca_major_AZ_PAF.pdf> [Accessed on

Edited by Alison Clements, Margaret David, Dong Lee, and Jacob Krebs, July 5, 2012

Jun 5, 2012].

Miller, J. H. 2003. Nonnative invasive plants of southern forests: A field guide for identification and control. Gen. Tech. Rep. SRS-62. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 42 pp. <http://www.srs.fs.usda.gov/pubs/gtr/gtr_srs062/> [Accessed on Jun 5, 2012].

Randall, J. M. and Marinelli, J. eds. 1996. Invasive plants: Weeds of the global garden. Handbook #149. Brooklyn, NY: Brooklyn Botanic Garden. 111 pp.

Stone, K. R. 2009. Vinca major, V. minor. In: Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). <<http://www.fs.fed.us/database/feis/>> [Accessed on Jun 5, 2012].

Swearingen, J., K. Reshetiloff., B. Slattery., and S. Zwicker S. 2002. Plant Invaders of Mid-Atlantic Natural Areas. National Park Service and U.S. Fish & Wildlife Service, 82 pp.

Weber, E. 2003. Invasive plant species of the world: a reference guide to environmental weeds. CABI Publishing, Cambridge, Massachusetts. 548 pp.

Glossary

Coverage. Visual or quantitative estimate of the relative amount of area in a stratum where the canopy of the non-native species intercepts the light that would otherwise be available for other species in or below that stratum. Estimated cover may be dispersed or continuous in a site. Cover is usually measured when foliage is fully expanded. In the case of species that form a dense, continuous mat of rhizomes or stolons, the percent of the soil surface or upper level occupied by that root mat can be estimated as soil, rather than canopy, cover.

Discrete sites. Disjunct habitat-types or fragments of habitats at least 1 mile apart that support invasive plant populations that likely arose by separate long-distance dispersal mechanisms.

Federal- or Indiana -listed. Species that are listed by Federal laws or Indiana statutes or rules as threatened or endangered within the State of Indiana. This list with notes is available at <http://www.state.in.us/dnr/naturepr/endanger/plant.htm>

Invasive. A species that forms self-sustaining and expanding populations within a natural plant community with which it had not previously been associated (Vitousek *et al.* 1995).

Long-term alterations in ecosystem processes. Examples of ecosystem processes that could be altered: erosion and sedimentation rates; land elevation; water channels; water-holding capacity; water-table depth; surface flow patterns; rates of nutrient mineralization or immobilization; soil or water chemistry; and type, frequency, intensity, or duration of disturbance. For further explanation see Gordon (1998).

Native. Species within its natural range or natural zone of dispersal (i.e., within the range it could have, or would have, occupied without direct or indirect introduction and/or care by humans. Excludes species descended from domesticated ancestors) (Vitousek *et al.* 1995).

Natural areas. Natural areas: Areas with native plant communities supporting native plant and animal species, with long undisturbed soil systems, and hydrological regimes relatively intact or under restoration. Edges of historically or currently disturbed areas (roadsides, trails, adjacent to historically disturbed locations, etc.) should not be included in the assessment of invasion into natural areas. That invasion may have been facilitated by the edges, but has to have extended into the native communities for inclusion in this category.